

11-29-00

A

PTO/SB/29 (12/97)

Approved for use through 09/30/00. OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

11/29/00
JCS67 U.S. PTO**UTILITY
PATENT APPLICATION
TRANSMITTAL**

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No.	7853-220	Total Pages	25
First Named Inventor or Application Identifier			
Moore et al.			
Express Mail Label No.	EL 501 636 208 US		

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner
Box Patent Application
Washington, DC 20231

1. ☒ Fee Transmittal Form
Submit an original, and a duplicate for fee processing

2. ☒ Specification
(preferred arrangement set forth below) [Total Pages 111]

-Descriptive title of the invention
-Cross Reference to Related Applications
-Statement Regarding Fed sponsored R&D
-Reference to Microfiche Appendix
-Background of the Invention
-Brief Summary of the Invention
-Brief Description of the Drawings *(if filed)*
-Detailed Description of the Invention *(including drawings, if filed)*
-Claim(s)
-Abstract of the Disclosure
☒ Drawing(s) *(35 USC 113)* [Total Sheets 6]

- ☒ Oath or Declaration *(unexecuted)* [Total Sheets 2]

- a. ☐ Newly executed (original or copy)
b. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 17 completed)
[Note Box 5 below]

1. ☐ **DELETION OF INVENTOR(S)**

Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33 (b).

- ☒ Incorporation By Reference *(useable if Box 4b is checked)*
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program *(Appendix)*
7. ☒ Nucleotide and/or Amino Acid Sequence Submission
(if applicable, all necessary)
a. ☐ Computer Readable Copy
b. ☒ Paper Copy *(identical to computer copy)*
c. ☒ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & document(s))
9. ☐ 37 CFR 3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)
10. ☐ English Translation Document *(if applicable)*
11. ☒ Information Disclosure ☐ Copies of IDS
Statement (IDS)/PTO-1449 ☐ Citations
12. ☒ Preliminary Amendment
13. ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
14. ☐ Small Entity ☐ Statement filed in prior application,
Statement(s) Status still proper and desired
15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)
16. ☐ Other:

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

☐ Continuation ☒ Divisional ☐ Continuation-in-part (CIP) of prior application No: 09/062,753 filed April 17, 1998

18. CORRESPONDENCE ADDRESS

☒ Customer Number or Bar Code Label 20583 *(Insert Customer No. or Attach bar code label here)* ☐ Correspondence address below

NAME			
ADDRESS			
CITY	STATE	ZIP CODE	
COUNTRY	TELEPHONE	FAX	

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

Express Mail No.: EL 501 636 208 US

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Prior application: Examiner Murphy, J.

Art Unit 1644

Assistant Commissioner for Patents
Box PATENT APPLICATION
Washington, D.C. 20231

Sir:

This is a request for filing a ☐ continuation ☒ divisional application under 37 CFR § 1.53(b), of pending prior application no. 09/062,753 filed on April 17, 1998.

of Karen Moore; Deborah Lynn Nagle; and Elizabeth A. Woolf
(inventor(s) currently of record in prior application)

for NOVEL HUMAN G-PROTEIN COUPLED RECEPTOR
(title of invention)

1. ☒ The filing fee is calculated below:

PATENT APPLICATION FEE VALUE

TYPE	NO. FILED	LESS	EXTRA	EXTRA RATE	FEE
Total Claims	73	-20	53	\$18.00 each	\$ 954.00
Independent	11	-3	8	\$80.00 each	\$ 640.00
Basic Fee					\$ 710.00
Multiple Dependency Fee If Applicable (\$270.00)					\$ 0.00
Total					\$ 2,304.00
50% Reduction for Independent Inventor, Nonprofit Organization or Small Business Concern					- \$ 0.00
Total Filing Fee					\$ 2,304.00

2. ☒ Please enter and consider the amendments to the claims made in the accompanying Preliminary Amendment before calculating the filing fee. Please charge the required fee to Pennie & Edmonds LLP Deposit Account No. 16-1150. A copy of this sheet is enclosed.
3. ☒ Amend the specification as follows: On Page 1, line 5, after "This application is a", insert -- divisional application of Serial No. 09/062,753, filed April 17, 1998, which is a --. On page 1, line 6, after "April 17, 1997," insert -- each of --.

- 4a. ☐ Transfer the drawings from the prior application to this application and abandon the prior application as of the filing date accorded this application. A duplicate copy of this sheet is enclosed for filing in the prior application file.
- 4b. ☐ New formal drawings are enclosed.
- 4c. ☒ Informal drawings are enclosed.
- 5a. ☐ Priority of application no. filed on in is claimed under 35 U.S.C. §119.
- 5b. ☐ The certified copy has been filed in prior application no. , filed .
6. ☐ The prior application is assigned of record to .
7. ☒ Unexecuted Declaration and Power of Attorney is enclosed herewith.
8. ☒ This application contains nucleic acid and/or amino acid sequences required to be disclosed in a Sequence Listing under 37 CFR §§1.821-1.825. It is requested that the Sequence Listing in computer readable form from prior application Serial No. 09/062,753, filed on April 17, 1998 be made a part of the present application as provided for by 37 C.F.R. §1.821(e). The sequences disclosed therein are the same as the sequences disclosed in this application. A copy of the paper Sequence Listing from application Serial No. 09/062,753 is enclosed.
9. ☒ The undersigned states, under 37 C.F.R. §1.821(f), that the content of the enclosed paper Sequence Listing from application Serial No. 09/062,753 is the same as the content of the computer readable form submitted in application Serial No. 09/062,753.
10. ☒ Additional enclosures or instructions. A copy of the following documents are enclosed: (a) an Information Disclosure Statement; (b) a revised PTO-1449 form; (c) a Preliminary Amendment; and (d) an Exhibit A (pending claims).

Respectfully submitted,

By: *Jennifer Chioda*
Reg No. 46,617

Date November 28, 2000

Laura A. Coruzzi 30,742

Laura A. Coruzzi (Reg. No.)
PENNIE & EDMONDS LLP
1155 Avenue of the Americas
New York, N.Y. 10036-2711
(212) 790-9090

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: Moore et al.

Application No.: To Be Assigned

Group Art Unit: To Be Assigned

Filed: Herewith

Examiner: To Be Assigned

For: A NOVEL HUMAN G-PROTEIN
COUPLED RECEPTOR

Attorney Docket No.: 7853-220

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

In connection with the above-identified application, and in accordance with the Rules of practice, please enter the following amendments and remarks.

IN THE CLAIMS:

Please cancel the Claims as follows:

Cancel Claims 1-15, without prejudice.

Amend Claims 16, 17, 52 and 60, as follows:

16. (amended) An antibody that immunospecifically binds [the gene product of Claim 14] to a polypeptide comprising:

- a. the amino acid sequence in SEQ ID NO:2 or 4;
- b. the amino acid sequence encoded by the nucleic acid insert of the clone contained in ATCC accession No. 98414; or
- c. the amino acid sequence encoded by the nucleic acid sequence in SEQ ID NO:1 or 3.

17. (amended) An antibody that immunospecifically binds [the gene product of Claim 15] a gene product encoded by a nucleic acid molecule which hybridizes under stringent conditions to the complement of:

- a. the nucleic acid molecule encoding a polypeptide comprising the amino acid sequence in SEQ ID NO:2 or 4;
- b. the nucleic acid insert of the clone contained in ATCC accession No. 98414; or
- c. the nucleic acid sequence in SEQ ID NO:1 or 3.

52. (amended) The method of Claim 51 wherein the compound comprises: [the nucleic acid molecule of Claim 1, 3 or 7.]:

- a. an isolated nucleic acid molecule encoding a polypeptide comprising the amino acid sequence in SEQ ID NO:2 or 4;
- b. the nucleic acid insert of the clone contained in ATCC accession No. 98414;
- c. an isolated nucleic acid molecule which hybridizes to the complement of the nucleic acid molecule encoding a polypeptide comprising the amino acid sequence in SEQ ID NO:2 or 4; or
- d. an isolated nucleic acid molecule comprising the nucleic acid sequence in SEQ ID NO:1 or 3.

60. (amended) The method of Claim 59 wherein the compound comprises [the nucleic acid molecule of Claims 1, 3, or 7.]:

- a. an isolated nucleic acid molecule encoding a polypeptide comprising the amino acid sequence in SEQ ID NO:2 or 4;
- b. the nucleic acid insert of the clone contained in ATCC accession No. 98414;
- c. an isolated nucleic acid molecule which hybridizes to the complement of the nucleic acid molecule encoding a polypeptide comprising the amino acid sequence in SEQ ID NO:2 or 4; or
- d. an isolated nucleic acid molecule comprising the nucleic acid sequence in SEQ ID NO:1 or 3.

IN THE SPECIFICATION:

Please amend the specification as follows:

At page 4, line 15, after "encoding human I5E", please insert the phrase

-- (SEQ ID NOS:1-2) --.

At page 4, line 17, after "encoding mouse I5E", please insert the phrase

-- (SEQ ID NOS:3-4) --.

At page 77, line 17, after "...GTCGG-3'", please insert the phrase

-- (SEQ ID NO:5) --.

At page 77, line 18, after "...TACCG-3'", please insert the phrase

-- (SEQ ID NO:6) --.

At page 79, line 36, after "...AGATG-3'", please insert the phrase

-- (SEQ ID NO:7) --.

At page 80, line 1, after "...CACTA-3'", please insert the phrase

-- (SEQ ID NO:8) --.

REMARKS

Applicants have cancelled Claims 1-15, without prejudice to Applicants' rights to pursue the subject matter of the claims in related applications. Applicants have amended Claims 16, 17, 52 and 60 to more particularly point out and distinctly claim the subject matter of the invention. The amendments to the claims are fully supported by the instant specification, see, *e.g.*, page 25, line 1 to page 63, line 23, and do not constitute new matter.

The specification has been amended pursuant to 37 C.F.R. § 1.821(d) to incorporate the sequence identifiers corresponding to the Sequence Listing submitted concurrently herewith. The incorporation of the sequence identifiers introduces no new matter.

[illegible]

7

Laura A. Coruzzi 30,742
Laura A. Coruzzi (Reg. No.)

Enclosures

William C. Kuyper
Reg. No. 39,201

A NOVEL HUMAN G-PROTEIN COUPLED RECEPTOR

5 This application is a Continuation-In-Part of Serial No. 08/833,226, filed on April 17, 1997, which is incorporated herein by reference in its entirety.

1. INTRODUCTION

0721302 113600
The present invention relates to the discovery, identification and characterization of nucleic acids that encode a novel G protein coupled receptor (referred to herein as I5E). The invention encompasses I5E nucleotides, host cell expression systems, I5E proteins, fusion proteins, polypeptides and peptides, antibodies to the receptor, transgenic animals that express a I5E transgene, or recombinant knock-out animals that do not express the I5E, antagonists and agonists of the receptor, and other compounds that modulate I5E gene expression or I5E activity that can be used for diagnosis, drug screening, clinical trial monitoring, and/or used to treat disorders such as inflammatory, central nervous system or metabolic disorders such as body weight disorders including obesity, cachexia and anorexia.

2. BACKGROUND OF THE INVENTION

25 Many biological processes are mediated by proteins participating in signal transduction pathways that involve G-proteins and/or second messengers. G-protein coupled receptors are plasma membrane proteins capable of transducing signals across a cell membrane so as to initiate a second messenger response. To this end, the G-protein coupled receptors bind a variety of ligands ranging from small biogenic amines to peptides, small proteins and large

been postulated to span the plasma membrane, connected by hydrophilic extracellular and intracellular loops. The G-protein family of coupled receptors includes dopamine, calcitonin, adrenergic, endothelia, CAMP, adenosine, 5 serotonin, follicle stimulating hormone, opsin and rhodopsin receptors.

G-protein coupled receptors can be intracellularly coupled to various intracellular enzymes, ion channels and transporters. Different G-protein α -subunits preferentially 10 stimulate particular effectors to modulate various biological functions in a cell. Phosphorylation of cytoplasmic residues of G-protein coupled receptors have been identified as an important mechanism for the regulation of G-protein coupling of the G-protein coupled receptors.

15

3. SUMMARY OF THE INVENTION

The present invention relates to the discovery, identification and characterization of nucleic acids that encode I5E, a novel G-protein coupled receptor protein that 20 contains regions of homology to the neuropeptide (NPY) receptor.

The invention encompasses the following nucleotides, host cells expressing such nucleotides, and the expression products of such nucleotides: (a) nucleotides that encode 25 mammalian I5Es, including the human I5E, and the I5E gene product; (b) nucleotides that encode portions of the I5E that correspond to its functional domains, and the polypeptide products specified by such nucleotide sequences, including but not limited to the extracellular domains (ECDs), the 30 transmembrane domains (TMs), and the cytoplasmic domains (CDs); (c) nucleotides that encode mutants of the I5E in which all or a part of one of the domains is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including but not limited to soluble 35 receptors in which one or more of the TM domains are deleted, and nonfunctional receptors in which all or a portion of the CD is deleted; (d) nucleotides that encode fusion proteins

containing the I5E or one of its domains (e.g., the extracellular domains) fused to another polypeptide.

The invention also encompasses agonists and antagonists of I5E, including small molecules, large molecules, mutant
5 natural I5E ligand proteins that compete with native natural I5E ligand, and antibodies, as well as nucleotide sequences that can be used to inhibit I5E gene expression (e.g., antisense and ribozyme molecules, and gene or regulatory sequence replacement constructs) or to enhance I5E gene
10 expression (e.g., expression constructs that place the I5E gene under the control of a strong promoter system), and transgenic animals that express an I5E transgene or "knock-outs" that do not express I5E.

Further, the present invention also relates to methods
15 for the use of the I5E gene and/or I5E gene products for the identification of compounds which modulate, i.e., act as agonists or antagonists, of I5E gene expression and or I5E gene product activity. In addition, the invention relates to methods of identifying compounds suitable for treatment of
20 diseases characterized by aberrant expression or activity levels of I5E. Such compounds can be used as therapeutic agents for use in treatment of immune disorders such as inflammation, central nervous system disorders, or metabolic disorders such as obesity, cachexia and anorexia. The
25 invention further relates to methods of treating diseases characterized by aberrant expression or activity of I5E. The methods of treatment involve the administration of compounds that act to modulate I5E expression or I5E activity.

3.1. DEFINITIONS

As used herein, the following terms, whether used in the singular or plural, will have the meanings indicated:

I5E nucleotides or coding sequences: means nucleotide sequences encoding I5E protein, polypeptide or peptide
35 fragments of I5E protein, or I5E fusion proteins. I5E nucleotide sequences encompass DNA, including genomic DNA (e.g. the I5E gene) or cDNA, or RNA.

I5E: means natural I5E ligand receptor protein.
Polypeptides or peptide fragments of I5E protein are referred to as I5E polypeptides or I5E peptides.
Fusions of I5E, or I5E polypeptides or peptide fragments to an unrelated protein are referred to herein as I5E fusion proteins.
A functional I5E refers to a protein which binds natural I5E ligand with high affinity in vivo or in vitro.

ECD: means "extracellular domain".

TM: means "transmembrane domain".

CD: means "cytoplasmic domain".

4. DESCRIPTION OF THE FIGURES

FIG. 1. Nucleotide sequence and deduced amino acid sequence of human I5E cDNA encoding human I5E.

FIG. 2. Nucleotide sequence and deduced amino acid sequence of mouse I5E cDNA encoding mouse I5E.

5. DETAILED DESCRIPTION OF THE INVENTION

I5E, described for the first time herein, is a novel G-protein coupled receptor protein, the human embodiment of which shares about 24% homology with the neuropeptide Y receptor (NPY-2 receptor) at the amino acid level. The novel I5E has been characterized as having seven hydrophobic domains which span the plasma membrane and are connected by hydrophilic extracellular and intracellular loops. In most cases, the stimulation of these receptors accelerates the turnover of phosphoinositides, with an amplitude that depends on the tissue in which the receptor is expressed.

The invention encompasses the use of I5E nucleotides, I5E proteins and peptides, as well as antibodies to I5E (which can, for example, act as I5E agonists or antagonists), antagonists that inhibit ligand binding, receptor activity or expression, or agonists that increase the binding affinity of the I5E ligand, activate receptor activity, or allow ligand to bind better or increase its expression in the diagnosis and treatment of disorders, including, but not limited to

09724392.112800

treatment of inflammatory, immune, central nervous system and metabolic disorders such as body weight disorders including obesity, cachexia and anorexia. The diagnosis of an I5E abnormality in a patient, or an abnormality in the I5E signal transduction pathway, will assist in devising a proper treatment or therapeutic regimen. In addition, I5E nucleotides and I5E proteins are useful for the identification of compounds effective in the treatment of disorders based on the aberrant expression or activity of I5E.

In particular, the invention described in the subsections below encompasses I5E, polypeptides or peptides corresponding to functional domains of the I5E (e.g., ECD, TM or CD), mutated, truncated or deleted I5Es (e.g. an I5E with one or more functional domains or portions thereof deleted, such as Δ TM and/or Δ CD), I5E fusion proteins (e.g. an I5E or a functional domain of I5E, such as one or more of the ECDs, fused to an unrelated protein or peptide such as an immunoglobulin constant region, i.e., IgFc), nucleotide sequences encoding such products, and host cell expression systems that can produce such I5E products.

The invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and agonists of the I5E, as well as compounds or nucleotide constructs that inhibit expression of the I5E gene (transcription factor inhibitors, antisense and ribozyme molecules, or gene or regulatory sequence replacement constructs), or promote expression of I5E (e.g., expression constructs in which I5E coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.). The invention also relates to host cells and animals genetically engineered to express the human I5E (or mutants thereof) or to inhibit or "knock-out" expression of the animal's endogenous I5E.

The I5E proteins or peptides, I5E fusion proteins, I5E nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant I5Es or

inappropriately expressed I5Es for the diagnosis of disorders including immune disorders such as inflammation, central nervous system disorders, and metabolic disorders such as body weight disorders including obesity, cachexia and anorexia. The I5E proteins or peptides, I5E fusion proteins, I5E nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs effective in the treatment of disorders. The use of engineered host cells and/or animals may offer an advantage in that such systems allow not only for the identification of compounds that bind to the ECD of the I5E, but can also identify compounds that affect the signal transduced by the activated I5E.

Finally, the I5E protein products (especially soluble derivatives such as peptides corresponding to the I5E ECD, or truncated polypeptides lacking the TM or CD domains) and fusion protein products (especially I5E-Ig fusion proteins, i.e., fusions of the I5E or a domain of the I5E, e.g., one or more ECDs, Δ TM to an IgFc), antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate signal transduction which may act on downstream targets in the I5E signal transduction pathway) can be used for therapy of such diseases. For example, the administration of an effective amount of soluble I5E ECD, Δ TM I5E or an ECD-IgFc fusion protein, or an anti-idiotypic antibody (or its Fab) that mimics the I5E ECD would "mop up" or "neutralize" endogenous ligand, and prevent or reduce binding and receptor activation. Nucleotide constructs encoding such I5E products can be used to genetically engineer host cells to express such I5E products in vivo; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of the I5E, I5E peptide, soluble ECD or Δ TM or I5E fusion protein that will "mop up" or neutralize natural I5E ligand. Nucleotide constructs encoding functional I5Es, mutant I5Es, as well as antisense and ribozyme molecules can be used in

"gene therapy" approaches for the modulation of I5E expression and/or activity in the treatment of disorders arising from the aberrant activity of the I5E.

5

5.1. THE I5E GENE

- The cDNA sequence (SEQ. ID. No. 1) and deduced amino acid sequence (SEQ. ID. No. 2) of human I5E are shown in FIG. 1. The cDNA sequence (SEQ. ID. No. 3) and deduced amino acid sequence (SEQ. ID. No. 4) of mouse I5E are shown in FIG.
- 10 2. The I5E nucleotide sequences of the invention include:
- (a) the DNA sequence shown in FIG. 1, FIG. 2, or contained in the cDNA clone as deposited with the American Type Culture Collection (ATCC) and assigned Accession No. 98414
- (b) nucleotide sequences that encode the amino acid sequence
- 15 shown in FIG. 1, FIG. 2, or contained in the cDNA clone as deposited with the American Type Culture Collection (ATCC) and assigned Accession No. 98414 (c) any nucleotide sequence that hybridizes to the complement of the DNA sequence shown in FIG. 1 or contained in the cDNA clone as deposited with
- 20 the American Type Culture Collection (ATCC) and assigned Accession No. 98414 under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989,
- 25 Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3); (d) any nucleotide sequence that hybridizes to the complement of the DNA sequence shown in FIG. 1 or continued in the cDNA clone as deposited with the
- 30 American Type Culture Collection (ATCC) and assigned Accession No. 98414 under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68° and encodes a functionally equivalent
- 35 gene product; and (e) any nucleotide sequence that hybridizes to the complement of the DNA sequences that encode the amino acid sequence shown in FIG. 1, FIG. 2, or contained in cDNA

clone as deposited with the ATCC and assigned Accession No. 98414, under less stringent conditions, such as moderately stringent conditions, e.g., washing in 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989, supra), and encodes a

5 functionally equivalent I5E gene product. Functional equivalents of the I5E include naturally occurring I5E present in other species, and mutant I5Es whether naturally occurring or engineered. The invention also includes degenerate variants of sequences (a) through (e).

10 Preferred I5E nucleic acids encode polypeptides that are at least 55% identical or similar to the amino acid sequence shown in FIG. 1 or the I5E amino acid sequence encoded by the cDNA clone as deposited with the ATCC and assigned Accession No. 98414. Nucleic acids which encode polypeptides which are
15 at least about 70%, and even more preferably at least about 80%, 85%, 90%, 95%, or 98% identical or similar with the amino acid sequence represented in FIG. 1 or the I5E amino acid sequence encoded by the cDNA clone as deposited with the ATCC and assigned Accession No. 98414 are also within the
20 scope of the invention. In a particularly preferred embodiment, the nucleic acid of the present invention encodes a polypeptide having an overall amino acid sequence homology or identity of at least about 70%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at
25 least about 98%, or at least 99% with the amino acid sequence shown in FIG. 1 or the I5E amino acid sequence encoded by the cDNA clone as deposited with the ATCC and assigned Accession No. 98414.

The invention also includes nucleic acid molecules,
30 preferably DNA molecules, that hybridize to, and are therefore the complements of, the nucleotide sequences (a) through (d), in the preceding paragraph. Such hybridization conditions may be highly stringent or less highly stringent, as described above. In instances wherein
35 the nucleic acid molecules are deoxyoligonucleotides ("oligos"), highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for

- 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as I5E antisense molecules, useful, for example, in I5E gene regulation (for and/or as antisense primers in amplification reactions of I5E gene nucleic acid sequences). With respect to I5E gene regulation, such techniques can be used to regulate, for example, inflammatory disease, treatment of pain, central nervous system disorders or gastrointestinal disorders.
- 10 Further, such sequences may be used as part of ribozyme and/or triple helix sequences, also useful for I5E gene regulation.

- In addition to the I5E nucleotide sequences described above, full length I5E cDNA or gene sequences present in the same species and/or homologs of the I5E gene present in other species can be identified and readily isolated, without undue experimentation, by molecular biological techniques well known in the art. The identification of homologs of I5E in related species can be useful for developing animal model systems more closely related to humans for purposes of drug discovery. For example, expression libraries of cDNAs synthesized from mRNA derived from the organism of interest can be screened using labeled natural I5E ligand derived from that species, e.g., a natural I5E ligand fusion protein.
- 25 Alternatively, such cDNA libraries, or genomic DNA libraries derived from the organism of interest can be screened by hybridization using the nucleotides described herein as hybridization or amplification probes. Furthermore, genes at other genetic loci within the genome that encode proteins which have extensive homology to one or more domains of the I5E gene product can also be identified via similar techniques. In the case of cDNA libraries, such screening techniques can identify clones derived from alternatively spliced transcripts in the same or different species.
- 30 Screening can be by filter hybridization, using duplicate filters. The labeled probe can contain at least 15-30 base pairs of the I5E nucleotide sequence, as shown in

FIG. 1, or FIG. 2. The hybridization washing conditions used should be of a lower stringency when the cDNA library is derived from an organism different from the type of organism from which the labeled sequence was derived. With respect to 5 the cloning of a I5E homolog, using human I5E probes, for example, hybridization can, for example, be performed at 65°C overnight in Church's buffer (7% SDS, 250 mM NaHPO₄, 2μM EDTA, 1% BSA). Washes can be done with 2XSSC, 0.1% SDS at 65°C and then at 0.1XSSC, 0.1% SDS at 65°C.

10 Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular 15 Cloning, A Laboratory Manual, Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y.

Alternatively, the labeled I5E nucleotide probe may be 20 used to screen a genomic library derived from the organism of interest, again, using appropriately stringent conditions. The identification and characterization of human genomic clones is helpful for designing diagnostic tests and clinical protocols for treating disorders in human patients. For 25 example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g. splice acceptor and/or donor sites), etc., that can be used 30 in diagnostics.

Further, an I5E gene homolog may be isolated from nucleic acid of the organism of interest by performing PCR using two degenerate oligonucleotide primer pools designed on the basis of amino acid sequences within the I5E gene product 35 disclosed herein. The template for the reaction may be cDNA obtained by reverse transcription of mRNA prepared from cell

08724392.112000

lines or tissue known or suspected to express an I5E gene allele.

The PCR product may be subcloned and sequenced to ensure that the amplified sequences represent the sequences of an I5E gene. The PCR fragment may then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment may be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment may be used to isolate genomic clones via the screening of a genomic library.

PCR technology may also be utilized to isolate full length cDNA sequences. For example, RNA may be isolated, following standard procedures, from an appropriate cellular or tissue source (i.e., one known, or suspected, to express the I5E gene). A reverse transcription reaction may be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNAase H, and second strand synthesis may then be primed with a poly-C primer. Thus, cDNA sequences upstream of the amplified fragment may easily be isolated. For a review of cloning strategies which may be used, see e.g., Sambrook et al., 1989, supra.

The I5E gene sequences may additionally be used to isolate mutant I5E gene alleles. Such mutant alleles may be isolated from individuals either known or proposed to have a genotype which contributes to the symptoms of disorders arising from the aberrant expression or activity of the I5E protein. Mutant alleles and mutant allele products may then be utilized in the therapeutic and diagnostic systems described below. Additionally, such I5E gene sequences can be used to detect I5E gene regulatory (e.g., promoter or promotor/enhancer) defects which can affect the expression of the I5E.

5 5' oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying the mutant I5E allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that 10 hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the 15 mutant I5E allele to that of the normal I5E allele, the mutation(s) responsible for the loss or alteration of function of the mutant I5E gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known 20 to carry the mutant I5E allele, or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express the mutant I5E allele. The normal I5E gene or any suitable fragment thereof may then be labeled and used as a probe to identify the corresponding mutant I5E allele in such 25 libraries. Clones containing the mutant I5E gene sequences may then be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated 30 from a tissue known, or suspected, to express a mutant I5E allele in an individual suspected of or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with 35 antibodies raised against the normal I5E gene product, as described, below, in Section 5.3. (For screening techniques, see, for example, WP, E. and Lane, eds., 1988, "Antibodies: A

000212632.112600

Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.) Additionally, screening can be accomplished by screening with labeled natural I5E ligand fusion proteins, such as, for example, AP-natural I5E ligand or natural I5E 5 ligand-AP fusion proteins. In cases where an I5E mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a frameshift mutation), a polyclonal set of antibodies to I5E are likely to cross-react with the mutant I5E gene product. Library clones detected 10 via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

The invention also encompasses nucleotide sequences that encode mutant I5Es, peptide fragments of the I5E, truncated 15 I5Es, and I5E fusion proteins. These include, but are not limited to nucleotide sequences encoding mutant I5Es described in section 5.2 infra; polypeptides or peptides corresponding to one or more of the ECDs, or TM and/or CD domains of the I5E or portions of these domains; truncated 20 I5Es in which one or two of the domains is deleted, e.g., a soluble I5E lacking a TM domain, or both a TM and CD regions, or a truncated, nonfunctional I5E lacking all, or a portion of a CD region. Nucleotides encoding fusion proteins may include by are not limited to full length I5E, truncated I5E 25 or peptide fragments of I5E fused to an unrelated protein or peptide, such as for example, a transmembrane sequence, which anchors the I5E ECD to the cell membrane; an Ig Fc domain which increases the stability and half life of the resulting fusion protein (e.g., I5E-Ig) in the bloodstream; or an 30 enzyme, fluorescent protein, luminescent protein which can be used as a marker.

The invention also encompasses (a) DNA vectors that contain any of the foregoing I5E coding sequences and/or their complements (i.e., antisense); (b) DNA expression 35 vectors that contain any of the foregoing I5E coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences; and

(c) genetically engineered host cells that contain any of the foregoing ISE coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell. As used herein, regulatory elements include but are not limited to inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the cytomegalovirus hCMV immediate early gene, the early or late promoters of SV40 adenovirus, the lac system, the trp system, the TAC system, the TRC system, the major operator and promoter regions of phage A, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase, the promoters of acid phosphatase, and the promoters of the yeast α -mating factors.

5.2. ISE PROTEINS AND POLYPEPTIDES

FIG. 1 shows the amino acid sequence of the human I5E protein. The I5E amino acid sequences of the invention include the amino acid sequence shown in FIG. 1 (SEQ. ID. No:2) or the amino acid sequence encoded by DNA as deposited with the ATCC and assigned Accession No. 98414. Polypeptides which are at least about 70%, and even more preferably at least about 80%, 85%, 90%, 95% or 98% identical or similar with the amino acid sequence represented by FIG. 1 or the amino acid sequence encoded by the cDNA clone as deposited

00724300 112800

site-directed mutations of the I5E coding sequence can be engineered (using site-directed mutagenesis techniques well known to those skilled in the art) to generate mutant I5Es with increased function, e.g., higher binding affinity for 5 natural I5E ligand, and/or greater signalling capacity; or decreased function, e.g., lower binding affinity for natural I5E ligand, and/or decreased signal transduction capacity.

For example, regions of identity may be determined by alignment of human I5E (FIG. 1) with I5E homologs from other 10 species. Mutant I5Es can be engineered so that regions of identity are maintained, whereas the variable residues are altered, e.g., by deletion or insertion of an amino acid residue(s) or by substitution of one or more different amino acid residues. Conservative alterations at the variable 15 positions can be engineered in order to produce a mutant I5E that retains function; e.g., natural I5E ligand binding affinity or signal transduction capability or both. Non-conservative changes can be engineered at these variable positions to alter function, e.g., natural I5E ligand binding 20 affinity or signal transduction capability, or both. Alternatively, where alteration of function is desired, deletion or non-conservative alterations of the conserved regions can be engineered. For example, deletion or non-conservative alterations (substitutions or insertions) of the 25 CD, e.g., amino acid residues of human I5E, or amino acid residues, or portions of the CD, of the human I5E (FIG. 1). Non-conservative alterations to the ECD shown can be engineered to produce mutant I5Es with altered binding affinity for natural I5E ligand. The same mutation strategy 30 can also be used to design mutant I5Es based on the alignment of human I5E and I5E homologs from other species.

Other mutations to the I5E coding sequence can be made to generate I5Es that are better suited for expression, scale up, etc. in the host cells chosen. For example, cysteine 35 residues can be deleted or substituted with another amino acid in order to eliminate disulfide bridges; N-linked glycosylation sites can be altered or eliminated to achieve,

15E. The DNA sequence encoding one or more of the ECDs can be ligated together directly or via a linker oligonucleotide that encodes a peptide spacer. Such linkers may encode flexible, glycine-rich amino acid sequences thereby alluring the domains that are strung together to assume a conformation that can bind the natural I5E ligand. Alternatively, nucleotide sequences encoding individual domains within the ECD can be used to express I5E peptides. A variety of methods can be used to construct expression vectors containing the I5E nucleotide sequences as described in Section 5.1 and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. See, for example, the techniques described in Sambrook et al., 1989, supra, and Ausubel et al., 1989, supra. Alternatively, RNA capable of encoding I5E nucleotide sequences may be chemically synthesized using, for example, synthesizers. See, for example, the techniques described in "Oligonucleotide Synthesis", 1984, Gait, M.J. ed., IRL Press, Oxford, which is incorporated by reference herein in its entirety.

A variety of host-expression vector systems may be utilized to express the I5E nucleotide sequences of the invention. Where the I5E peptide or polypeptide is a soluble derivative (e.g., I5E peptides corresponding to the ECD; truncated or deleted I5E in which the TM and/or CD are deleted) the peptide or polypeptide can be recovered from the culture, i.e., from the host cell in cases where the I5E peptide or polypeptide is not secreted, and from the culture media in cases where the I5E peptide or polypeptide is secreted by the cells. However, the expression systems also encompass engineered host cells that express the I5E or functional equivalents in situ, i.e., anchored in the cell membrane. Purification or enrichment of the I5E from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well known to those skilled in the art. However, such engineered host cells

themselves may be used in situations where it is important not only to retain the structural and functional characteristics of the I5E, but to assess biological activity, e.g., in drug screening assays.

- 5 The expression systems that may be used for purposes of the invention include but are not limited to microorganisms such as bacteria (e.g., E. coli, B. subtilis) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing I5E nucleotide sequences; yeast
- 10 (e.g., Saccharomyces, Pichia) transformed with recombinant yeast expression vectors containing the I5E nucleotide sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the I5E sequences; plant cell systems infected with recombinant
- 15 virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing I5E nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant
- 20 expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

- In bacterial systems, a number of expression vectors may
- 25 be advantageously selected depending upon the use intended for the I5E gene product being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of I5E protein or for raising antibodies to the I5E protein, for example,
 - 30 vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the E. coli expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which the I5E coding sequence may be ligated
 - 35 individually into the vector in frame with the lacZ coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van

Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are
5 soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released
10 from the GST moiety.

In an insect system, Autographa californica nuclear polyhidrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in Spodoptera frugiperda cells. The I5E gene coding sequence may be cloned
15 individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of I5E gene coding sequence will result in inactivation of the polyhedrin gene and production of non-
20 occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect Spodoptera frugiperda cells in which the inserted gene is expressed. (E.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S.
25 Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the I5E nucleotide sequence of interest may be ligated to an
30 adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3)
35 will result in a recombinant virus that is viable and capable of expressing the I5E gene product in infected hosts. (E.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-

3659). Specific initiation signals may also be required for efficient translation of inserted I5E nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire I5E gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of the I5E coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (See Bittner et al., 1987, Methods in Enzymol. 153:516-544).

20 In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3 and WI38.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell

lines which stably express the I5E sequences described above may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the I5E gene product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the I5E gene product.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk⁻, hgprt⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hyg^r, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147).

Alternatively, any fusion protein may be readily purified by utilizing an antibody specific for the fusion

protein being expressed. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88: 8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺-nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

The I5E gene products can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate I5E transgenic animals.

Any technique known in the art may be used to introduce the I5E transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to pronuclear microinjection (Hoppe, P.C. and Wagner, T.E., 1989, U.S. Pat. No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci., USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

The present invention provides for transgenic animals that carry the I5E transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, i.e., mosaic animals. The transgene may be integrated as a single transgene or in concatamers, e.g., head-to-head

5.3. ANTIBODIES TO I5E PROTEINS

Antibodies that specifically recognize one or more epitopes of I5E, or epitopes of conserved variants of I5E, or peptide fragments of the I5E are also encompassed by the invention. Such antibodies include but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

The antibodies of the invention may be used, for example, in the detection of the I5E in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of I5E. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes, as described, below, in Section 5.5, for the evaluation of the effect of test compounds on expression and/or activity of the I5E gene product. Additionally, such antibodies can be used in conjunction with the gene therapy techniques described, below, in Section 5.6, to, for example, evaluate the normal and/or engineered I5E-expressing cells prior to their introduction into the patient. Such antibodies may additionally be used as a method for the inhibition of abnormal I5E activity.

For the production of antibodies, various host animals may be immunized by injection with the I5E, an I5E peptide (e.g., one corresponding to a functional domain of the receptor, such as ECD, TM or CD), truncated I5E polypeptides (I5E in which one or more domains, e.g., the TM or CD, has been deleted), functional equivalents of the I5E or mutants of the I5E. Such host animals may include but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as

lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and Corynebacterium parvum. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, may be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated in vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger et al., 1984, Nature, 312:604-608; Takeda et al., 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region.

In addition, techniques have been developed for the production of humanized antibodies. (See, et al., Queen, U.S. Patent No. 5,585,089, which is incorporated herein by reference in its entirety.) An immunoglobulin light or heavy

chain variable region consists of a "framework" region interrupted by three hypervariable regions, referred to as complementarily determining regions (CDRs). The extent of the framework region and CDRs have been precisely defined
5 (see, "Sequences of Proteins of Immunological Interest", Kabat, E. et al., U.S. Department of Health and Human Services (1983). Briefly, humanized antibodies are antibody molecules from non-human species having one or more CDRs from the non-human species and a framework region from a human
10 immunoglobulin molecule.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, Science 242:423-426; Huston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward et al., 1989, Nature
15 334:544-546) can be adapted to produce single chain antibodies against I5E gene products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

20 Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include but are not limited to: the $F(ab')_2$ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be
25 generated by reducing the disulfide bridges of the $F(ab')_2$ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

30 Antibodies to the I5E can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" the I5E, using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, 1993, FASEB J 7(5):437-444; and Nissinoff, 1991, J. Immunol. 147(8):2429-2438). For example
35 antibodies which bind to the I5E ECD and competitively inhibit the binding of natural I5E ligand to the I5E can be used to generate anti-idiotypes that "mimic" the ECD and,

therefore, bind and neutralize natural I5E ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize the physiological activity of the natural gpc-R ligand.

5

5.4. SCREENING ASSAYS FOR COMPOUNDS THAT MODULATE I5E EXPRESSION OR ACTIVITY

The following assays are designed to identify compounds that are capable of modulating the expression or biological activity of I5E. Such compounds can be used to treat diseases arising from aberrant expression or activity of the I5E. Such diseases include immune disorders such as inflammation, central nervous system disorders or metabolic disorders such as those involved in body weight disorders, including but not limited to obesity, cachexia and anorexia.

The following assays are designed to identify compounds that interact with (e.g., bind to) I5E (including, but not limited to the ECD or CD of I5E), compounds that interact with (e.g., bind to) intracellular proteins that interact with I5E (including, but not limited to, the TM and CD of I5E), compounds that interfere with the interaction of I5E with transmembrane or intracellular proteins involved in I5E-mediated signal transduction, and to compounds which modulate the activity of I5E gene (i.e., modulate the level of I5E gene expression) or modulate the level of I5E. Such assays may be used to identify compounds that function as antagonist or agonists of I5E activity. Assays may additionally be utilized which identify compounds which bind to I5E gene regulatory sequences (e.g., promoter sequences) and which may modulate I5E gene expression. See e.g., Platt, K.A., 1994, J. Biol. Chem. 269:28558-28562, which is incorporated herein by reference in its entirety.

The compounds which may be screened in accordance with the invention include, but are not limited to peptides, antibodies and fragments thereof, and other organic compounds (e.g., peptidomimetics) that interact with (i.e., bind to) the ECD of the I5E and either mimic the activity triggered by

the natural ligand (i.e., agonists) or inhibit the activity triggered by the natural ligand (i.e., antagonists); as well as peptides, antibodies or fragments thereof, and other organic compounds that mimic the ECD of the I5E (or a portion thereof) and bind to and "neutralize" natural ligand.

Such compounds may include, but are not limited to, peptides such as, for example, soluble peptides, including but not limited to members of random peptide libraries; (see, e.g., Lam, K.S. et al., 1991, Nature 354:82-84; Houghten, R. et al., 1991, Nature 354:84-86), and combinatorial chemistry-derived molecular library made of D- and/or L- configuration amino acids, phosphopeptides (including, but not limited to, members of random or partially degenerate, directed phosphopeptide libraries; see, e.g., Songyang, Z. et al., 1993, Cell 72:767-778); molecules from natural product libraries, antibodies (including, but not limited to, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab')₂ and FAb expression library fragments, and epitope-binding fragments thereof), and small organic or inorganic molecules.

Other compounds which can be screened in accordance with the invention include but are not limited to small organic molecules that are able to cross the blood-brain barrier, gain entry into an appropriate cell and affect the expression of the I5E gene or some other gene involved in the I5E signal transduction pathway (e.g., by interacting with the regulatory region or transcription factors involved in gene expression); or such compounds that affect the activity of the I5E or the activity of some other intracellular factor involved in the I5E signal transduction pathway.

5.4.1 ANIMAL-AND CELL-BASED MODEL SYSTEMS

Described herein are cell- and animal-based systems which act as models for disorders arising from aberrant expression or activity of I5E. Cell- and animal-based model systems can also be used to further characterize the activity of the I5E gene. Such assays can be utilized as part of

screening strategies designed to identify compounds which are capable of ameliorating I5E based disorders such as immune disorders, central nervous system disorders or metabolic disorders such as those involved in body weight disorders, including but not limited to obesity, cachexia and anorexia. Thus, the animal- and cell-based models can be used to identify drugs, pharmaceuticals, therapies and interventions which can be effective in treating disorders aberrant expression or activity of the I5E cytokine. In addition, as described in detail, below, in Section 5.7.1, such animal models can be used to determine the LD₅₀ and the ED₅₀ in animal subjects, and such data can be used to determine the *in vivo* efficacy of potential I5E disorder treatments.

Animal-based model systems of I5E based disorders such as, but not limited to, TH cell subpopulation-related disorders, based on aberrant I5E expression or activity, can include both non-recombinant animals as well as recombinantly engineered transgenic animals.

Animal models for I5E disorders can include, for example, genetic models. Animal models exhibiting I5E based disorder-like symptoms can be engineered by utilizing, for example, I5E sequences such as those described, above, in Section 5.2, in conjunction with techniques for producing transgenic animals that are well known to those of skill in the art. For example, I5E sequences can be introduced into, and overexpressed and/or misexpressed in, the genome of the animal of interest, or, if endogenous I5E sequences are present, they can either be overexpressed, misexpressed, or, alternatively, can be disrupted in order to underexpress or inactivate I5E gene expression.

In order to overexpress or misexpress a I5E gene sequence, the coding portion of the I5E gene sequence can be ligated to a regulatory sequence which is capable of driving high level gene expression or expression in a cell type in which the gene is not normally expressed in the animal and/or cell type of interest. Such regulatory regions will be well

known to those of skill in the art, and can be utilized in the absence of undue experimentation.

For underexpression of an endogenous I5E gene sequence, such a sequence can be isolated and engineered such that when reintroduced into the genome of the animal of interest, the endogenous I5E gene alleles will be inactivated, or "knocked-out". Preferably, the engineered I5E gene sequence is introduced via gene targeting such that the endogenous I5E sequence is disrupted upon integration of the engineered I5E gene sequence into the animal's genome. Gene targeting is discussed, below, in this Section.

Animals of any species, including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, and non-human primates, e.g., baboons, squirrels, monkeys, and chimpanzees can be used to generate animal models of I5E-related disorders.

Any technique known in the art can be used to introduce a I5E transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to pronuclear microinjection (Hoppe, P.C. and Wagner, T.E., 1989, U.S. Pat. No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci., USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

The present invention provides for transgenic animals that carry the I5E transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, i.e., mosaic animals. (See, for example, techniques described by Jakobovits, 1994, Curr. Biol. 4:761-763.) The transgene can be integrated as a single transgene or in concatamers, e.g., head-to-head tandems or head-to-tail

tandems. The transgene can also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko, M. et al., 1992, Proc. Natl. Acad. Sci. USA 89:6232-6236). The

- 5 regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

- When it is desired that the I5E transgene be integrated into the chromosomal site of the endogenous I5E gene, gene
10 targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous I5E gene of interest (e.g., nucleotide sequences of the mouse I5E gene, as depicted in FIG. 2) are designed for the purpose of integrating, via
15 homologous recombination with chromosomal sequences, into and disrupting the function of, the nucleotide sequence of the endogenous I5E gene. The transgene can also be selectively introduced into a particular cell type, thus inactivating the endogenous gene of interest in only that cell type, by
20 following, for example, the teaching of Gu et al. (Gu, H. et al., 1994, Science 265:103-106). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

- 25 Once transgenic animals have been generated, the expression of the recombinant I5E gene and protein can be assayed utilizing standard techniques. Initial screening can be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the
30 transgene has taken place. The level of mRNA expression of the I5E transgene in the tissues of the transgenic animals can also be assessed using techniques which include but are not limited to Northern blot analysis of tissue samples obtained from the animal, in situ hybridization analysis, and
35 RT-PCR. Samples of target gene-expressing tissue, can also be evaluated immunocytochemically using antibodies specific for the target gene transgene gene product of interest.

Further, the fingerprint pattern of gene expression of cells of interest can be analyzed and compared to the normal, non-I5E-related disorder fingerprint pattern. Those compounds which cause cells exhibiting I5E-related disorder-
5 like cellular phenotypes to produce a fingerprint pattern more closely resembling a normal fingerprint pattern for the cell of interest can be considered candidates for further testing regarding an ability to ameliorate I5E-related disorder symptoms.

- 10 In accordance with the invention, a cell-based assay system can be used to screen for compounds that modulate the activity of the I5E. To this end, cells that endogenously express I5E can be used to screen for compounds. Alternatively, cell lines, such as 293 cells, COS cells, CHO
15 cells, fibroblasts, and the like, genetically engineered to express the I5E can be used for screening purposes. Preferably, host cells genetically engineered to express a functional receptor that responds to activation by the natural I5E ligand can be used as an endpoint in the assay;
20 e.g., as measured by a chemical, physiological, biological, or phenotypic change, induction of a host cell gene or a reporter gene, change in cAMP levels, adenylyl cyclase activity, host cell G protein activity, extracellular acidification rate, host cell kinase activity, proliferation,
25 differentiation, etc.

To be useful in screening assays, the host cells expressing functional I5E should give a significant response to I5E ligand, preferably greater than 5-fold induction over background.

- 30 In utilizing such cell systems, the cells expressing the I5E are exposed to a test compound or to vehicle controls (e.g., placebos). After exposure, the cells can be assayed to measure the expression and/or activity of components of the signal transduction pathway of the I5E, or the activity
35 of the signal transduction pathway itself can be assayed. For example, after exposure, cell lysates can be assayed for induction of phospholipase C or accumulation of inositol

phosphate (IP) in the cell. The ability of a test compound to increase levels of phospholipase C or inositol phosphate, above those levels seen with cells treated with a vehicle control, indicates that the test compound induces signal transduction mediated by the I5E expressed by the host cell.

To determine intracellular inositol phosphate concentrations, an assay that utilizes [³H] inositol and anion exchange columns containing AG 1-X8 resin may be used as described in Tian et al. (1996, J.Neurochemistry 67:1191-1199). The assay may be performed in 96-well plates to enable high-throughput screening and 96 well-based scintillation counting instruments such as those manufactured by Wallac or Packard may be used for readout.

Other screening techniques include the use of cells which express the G-protein coupled receptor (for example, transfected CHO cells) in a system which measures extracellular pH changes caused by receptor activation, for example, as described in Science, volume 246, pages 181-296 (October 1989). In addition, a cytosensor microphysiometer can be used to detect and monitor the response of cells to chemical substances as described in McConnell et al. (• 1992, Science 257: 1906-1912). For example, potential agonists or antagonists may be contacted with a cell which expresses the G-protein coupled receptor and a second messenger response, e.g., signal transduction or pH changes, may be measured to determine whether the potential agonist or antagonist is effective.

Yet another screening procedure involves the use of the melanophores which are transfected to express the G-protein coupled receptor of the present invention. Such a screening technique is described in PCT WO 92/01810 published February 6, 1992.

Thus, for example, such an assay may be employed for screening for a receptor antagonist by contacting the melanophore cells which encode the G-protein coupled receptor with both the receptor ligand and a compound to be screened. Inhibition of the signal generated by the ligand indicates

that a compound is a potential antagonist for the receptor, i.e., inhibits activation of the receptor. The screen may also be employed for determining an agonist by contacting such cells with compounds to be screened and determining whether such compound generates a signal, i.e., activates the receptor.

Another such screening technique involves introducing RNA encoding I5E into *Xenopus* oocytes to transiently express the receptor. The receptor expressing oocytes may then be contacted, in the case of antagonist screening, with the receptor ligand and a compound to be screened, followed by detection of inhibition of a calcium signal.

Another method involves screening for antagonists by determining inhibition of binding of a labeled ligand to cells which have the receptor on the surface thereof. Such a method involves transfecting a eukaryotic cell with DNA encoding the G-protein coupled receptor such that the cell expresses the receptor on its surface and contacting the cell with a potential antagonist in the presence of a labeled form of a known ligand. The ligand can be labeled, e.g., by radioactivity. The amount of labeled ligand bound to the receptors is measured, e.g., by measuring radioactivity of the receptors. If the potential antagonist binds to the receptor as determined by a reduction of labeled ligand which binds to the receptors, the binding of labeled ligand to the receptor is inhibited.

Activation of G-protein coupled receptors normally results in induction of cAMP. Therefore, in an additional specific embodiment of the invention, constructs containing the cAMP responsive element linked to any of a variety of different reporter genes may be introduced into cells expressing the I5E receptor. Such reporter genes may include but is not limited to chloramphenicol acetyltransferase (CAT), luciferase, GUS, growth hormone, or placental alkaline phosphatase (SEAP). Following exposure of the cells to the test compound, the level of reporter gene expression may be quantitated to determine the test compound's ability to

regulate receptor activity. Alkaline phosphatase assays are particularly useful in the practice of the invention as the enzyme is secreted from the cell. Therefore, tissue culture supernatant may be assayed for secreted alkaline phosphatase.

- 5 In addition, alkaline phosphatase activity may be measured by calorimetric, bioluminescent or chemiluminescent assays such as those described in Bronstein, I. et al. (1994, Biotechniques 17: 172-177). Such assays provide a simple, sensitive easily automatable detection system for
- 10 pharmaceutical screening.

- Computer modelling and searching technologies permit identification of compounds, or the improvement of already identified compounds, that can modulate ISE expression or activity. Having identified such a compound or composition,
- 15 the active sites or regions are identified. Such active sites might typically be ligand binding sites, such as the interaction domains of natural ISE ligand with ISE itself. The active site can be identified using methods known in the art including, for example, from the amino acid sequences of
- 20 peptides, from the nucleotide sequences of nucleic acids, or from study of complexes of the relevant compound or composition with its natural ligand. In the latter case, chemical or X-ray crystallographic methods can be used to find the active site by finding where on the factor the
- 25 complexed ligand is found.

- Next, the three dimensional geometric structure of the active site is determined. This can be done by known methods, including X-ray crystallography, which can determine a complete molecular structure. On the other hand, solid or
- 30 liquid phase NMR can be used to determine certain intramolecular distances. Any other experimental method of structure determination can be used to obtain partial or complete geometric structures. The geometric structures may be measured with a complexed ligand, natural or artificial,
- 35 which may increase the accuracy of the active site structure determined.

If an incomplete or insufficiently accurate structure is determined, the methods of computer based numerical modelling can be used to complete the structure or improve its accuracy. Any recognized modelling method may be used, including parameterized models specific to particular biopolymers such as proteins or nucleic acids, molecular dynamics models based on computing molecular motions, statistical mechanics models based on thermal ensembles, or combined models. For most types of models, standard molecular force fields, representing the forces between constituent atoms and groups, are necessary, and can be selected from force fields known in physical chemistry. The incomplete or less accurate experimental structures can serve as constraints on the complete and more accurate structures computed by these modeling methods.

Finally, having determined the structure of the active site, either experimentally, by modeling, or by a combination, candidate modulating compounds can be identified by searching databases containing compounds along with information on their molecular structure. Such a search seeks compounds having structures that match the determined active site structure and that interact with the groups defining the active site. Such a search can be manual, but is preferably computer assisted. These compounds found from this search are potential ISE modulating compounds.

Alternatively, these methods can be used to identify improved modulating compounds from an already known modulating compound or ligand. The composition of the known compound can be modified and the structural effects of modification can be determined using the experimental and computer modelling methods described above applied to the new composition. The altered structure is then compared to the active site structure of the compound to determine if an improved fit or interaction results. In this manner systematic variations in composition, such as by varying side groups, can be quickly evaluated to obtain modified

modulating compounds or ligands of improved specificity or activity.

Further experimental and computer modeling methods useful to identify modulating compounds based upon
5 identification of the active sites of natural I5E ligand, I5E, and related transduction and transcription factors will be apparent to those of skill in the art.

Examples of molecular modelling systems are the CHARMM and QUANTA programs (Polygen Corporation, Waltham, MA).
10 CHARMM performs the energy minimization and molecular dynamics functions. QUANTA performs the construction, graphic modelling and analysis of molecular structure. QUANTA allows interactive construction, modification, visualization, and analysis of the behavior of molecules with
15 each other.

A number of articles review computer modelling of drugs interactive with specific proteins, such as Rotivinen, et al., 1988, Acta Pharmaceutical Fennica 97:159-166; Ripka, New Scientist 54-57 (June 16, 1988); McKinaly and Rossmann, 1989,
20 Annu. Rev. Pharmacol. Toxicol. 29:111-122; Perry and Davies, OSAR: Quantitative Structure-Activity Relationships in Drug Design pp. 189-193 (Alan R. Liss, Inc. 1989); Lewis and Dean, 1989 Proc. R. Soc. Lond. 236:125-140 and 141-162; and,
25 Askew, et al., 1989, J. Am. Chem. Soc. 111:1082-1090. Other computer programs that screen and graphically depict chemicals are available from companies such as BioDesign, Inc. (Pasadena, CA.), Allelix, Inc. (Mississauga, Ontario, Canada), and Hypercube, Inc. (Cambridge, Ontario). Although
30 these are primarily designed for application to drugs specific to particular proteins, they can be adapted to design of drugs specific to regions of DNA or RNA, once that region is identified.

Although described above with reference to design and
35 generation of compounds which could alter binding, one could also screen libraries of known compounds, including natural products or synthetic chemicals, and biologically active

materials, including proteins, for compounds which are inhibitors or activators.

Compounds identified via assays such as those described herein may be useful, for example, in elaborating the biological function of the I5E gene product, and for ameliorating disorders arising from the activity of the I5E.

5.4.2. IN VITRO SCREENING ASSAYS FOR COMPOUNDS THAT BIND TO I5E

In vitro systems may be designed to identify compounds capable of interacting with (e.g., binding to) I5E (including, but not limited to, the one or more ECDs or CDs of I5E). Compounds identified may be useful, for example, in modulating the activity of wild type and/or mutant I5E gene products; may be useful in elaborating the biological function of the I5E; may be utilized in screens for identifying compounds that disrupt normal I5E interactions; or may in themselves disrupt such interactions.

The principle of the assays used to identify compounds that bind to the I5E involves preparing a reaction mixture of the I5E and the test compound under conditions and for a time sufficient to allow the two components to interact and bind, thus forming a complex which can be removed and/or detected in the reaction mixture. The I5E species used can vary depending upon the goal of the screening assay. For example, where agonists of the natural ligand are sought, the full length I5E, or a soluble truncated I5E, e.g., in which the one or more of the TM and/or CD domains is deleted from the molecule, a peptide corresponding to one or more of the ECDs or TMs, or a fusion protein containing one or more of the I5E ECDs fused to a protein or polypeptide that affords advantages in the assay system (e.g., labeling, isolation of the resulting complex, etc.) can be utilized. Where compounds that interact with the cytoplasmic domain are sought to be identified, peptides corresponding to the I5E CD and fusion proteins containing the I5E CD can be used.

The screening assays can be conducted in a variety of ways. For example, one method to conduct such an assay would involve anchoring the I5E protein, polypeptide, peptide or fusion protein or the test substance onto a solid phase and detecting I5E/test compound complexes anchored on the solid phase at the end of the reaction. In one embodiment of such a method, the I5E reactant may be anchored onto a solid surface, and the test compound, which is not anchored, may be labeled, either directly or indirectly.

10 In practice, microtitre plates may conveniently be utilized as the solid phase. The anchored component may be immobilized by non-covalent or covalent attachments. Non-covalent attachment may be accomplished by simply coating the solid surface with a solution of the protein and drying.

15 Alternatively, an immobilized antibody, preferably a monoclonal antibody, specific for the protein to be immobilized may be used to anchor the protein to the solid surface. The surfaces may be prepared in advance and stored.

In order to conduct the assay, the nonimmobilized
20 component is added to the coated surface containing the anchored component. After the reaction is complete, unreacted components are removed (e.g., by washing) under conditions such that any complexes formed will remain immobilized on the solid surface. The detection of complexes
25 anchored on the solid surface can be accomplished in a number of ways. Where the previously nonimmobilized component is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the previously nonimmobilized component is not pre-labeled, an
30 indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the previously nonimmobilized component (the antibody, in turn, may be directly labeled or indirectly labeled with a labeled anti-Ig antibody).

35 Alternatively, a reaction can be conducted in a liquid phase, the reaction products separated from unreacted components, and complexes detected; e.g., using an

immobilized antibody specific for I5E protein, polypeptide, peptide or fusion protein or the test compound to anchor any complexes formed in solution, and a labeled antibody specific for the other component of the possible complex to detect
5 anchored complexes.

5.4.3. ASSAYS FOR INTRACELLULAR PROTEINS THAT INTERACT WITH THE I5E

Any method suitable for detecting protein-protein
10 interactions may be employed for identifying transmembrane proteins or intracellular proteins that interact with I5E. Among the traditional methods which may be employed are co-immunoprecipitation, crosslinking and co-purification through gradients or chromatographic columns of cell lysates
15 or proteins obtained from cell lysates and the I5E to identify proteins in the lysate that interact with the I5E. For these assays, the I5E component used can be a full length I5E, a soluble derivative lacking the membrane-anchoring region (e.g., a truncated I5E in which the TM is deleted
20 resulting in a truncated molecule containing the ECD fused to the CD), a peptide corresponding to the CD or a fusion protein containing the CD of I5E. Once isolated, such an intracellular protein can be identified and can, in turn, be used, in conjunction with standard techniques, to identify
25 proteins with which it interacts. For example, at least a portion of the amino acid sequence of an intracellular protein which interacts with the I5E can be ascertained using techniques well known to those of skill in the art, such as via the Edman degradation technique. (See, e.g., Creighton,
30 1983, "Proteins: Structures and Molecular Principles", W.H. Freeman & Co., N.Y., pp.34-49). The amino acid sequence obtained may be used as a guide for the generation of oligonucleotide mixtures that can be used to screen for gene sequences encoding such intracellular proteins. Screening
35 may be accomplished, for example, by standard hybridization or PCR techniques. Techniques for the generation of oligonucleotide mixtures and the screening are well-known.

(See, e.g., Ausubel, supra., and PCR Protocols: A Guide to Methods and Applications, 1990, Innis, M. et al., eds. Academic Press, Inc., New York).

Additionally, methods may be employed which result in
5 the simultaneous identification of genes which encode the transmembrane or intracellular proteins interacting with I5E. These methods include, for example, probing expression, libraries, in a manner similar to the well known technique of antibody probing of λ gt11 libraries, using labeled I5E
10 protein, or an I5E polypeptide, peptide or fusion protein, e.g., an I5E polypeptide or I5E domain fused to a marker (e.g., an enzyme, fluor, luminescent protein, or dye), or an Ig-Fc domain.

One method which detects protein interactions in vivo,
15 the yeast two-hybrid system, is described in detail for illustration only and not by way of limitation. One version of this system has been described (Chien et al., 1991, Proc. Natl. Acad. Sci. USA, 88:9578-9582) and is commercially available from Clontech (Palo Alto, CA).

20 Briefly, utilizing such a system, plasmids are constructed that encode two hybrid proteins: one plasmid consists of nucleotides encoding the DNA-binding domain of a transcription activator protein fused to an I5E nucleotide sequence encoding I5E, an I5E polypeptide, peptide or fusion
25 protein, and the other plasmid consists of nucleotides encoding the transcription activator protein's activation domain fused to a cDNA encoding an unknown protein which has been recombined into this plasmid as part of a cDNA library. The DNA-binding domain fusion plasmid and the cDNA library
30 are transformed into a strain of the yeast Saccharomyces cerevisiae that contains a reporter gene (e.g., HBS or lacZ) whose regulatory region contains the transcription activator's binding site. Either hybrid protein alone cannot activate transcription of the reporter gene: the DNA-binding
35 domain hybrid cannot because it does not provide activation function and the activation domain hybrid cannot because it cannot localize to the activator's binding sites.

Interaction of the two hybrid proteins reconstitutes the functional activator protein and results in expression of the reporter gene, which is detected by an assay for the reporter gene product.

- 5 The two-hybrid system or related methodology may be used to screen activation domain libraries for proteins that interact with the "bait" gene product. By way of example, and not by way of limitation, I5E may be used as the bait gene product. Total genomic or cDNA sequences are fused to
- 10 the DNA encoding an activation domain. This library and a plasmid encoding a hybrid of a bait I5E gene product fused to the DNA-binding domain are cotransformed into a yeast reporter strain, and the resulting transformants are screened for those that express the reporter gene. For example, and
- 15 not by way of limitation, a bait I5E gene sequence, such as the open reading frame of I5E (or a domain of I5E), as depicted in FIG. 1 can be cloned into a vector such that it is translationally fused to the DNA encoding the DNA-binding domain of the GAL4 protein. These colonies are purified and
- 20 the library plasmids responsible for reporter gene expression are isolated. DNA sequencing is then used to identify the proteins encoded by the library plasmids.

- A cDNA library of the cell line from which proteins that interact with bait I5E gene product are to be detected can be
- 25 made using methods routinely practiced in the art. According to the particular system described herein, for example, the cDNA fragments can be inserted into a vector such that they are translationally fused to the transcriptional activation domain of GAL4. This library can be co-transformed along
- 30 with the bait I5E gene-GAL4 fusion plasmid into a yeast strain which contains a lacZ gene driven by a promoter which contains GAL4 activation sequence. A cDNA encoded protein, fused to GAL4 transcriptional activation domain, that interacts with bait I5E gene product will reconstitute an
- 35 active GAL4 protein and thereby drive expression of the HIS3 gene. Colonies which express HIS3 can be detected by their growth on petri dishes containing semi-solid agar based media

lacking histidine. The cDNA can then be purified from these strains, and used to produce and isolate the bait I5E gene-interacting protein using techniques routinely practiced in the art.

5

**5.4.4. ASSAYS FOR COMPOUNDS THAT INTERFERE
WITH I5E/INTRACELLULAR OR I5E/
TRANSMEMBRANE MACROMOLECULE INTERACTION**

The macromolecules that interact with the I5E are referred to, for purposes of this discussion, as "binding
10 partners". These binding partners are likely to be involved in the I5E signal transduction pathway. Therefore, it is desirable to identify compounds that interfere with or disrupt the interaction of such binding partners with natural I5E ligand which may be useful in regulating the activity of
15 the I5E and control disorders associated with I5E activity.

The basic principle of the assay systems used to identify compounds that interfere with the interaction between the I5E and its binding partner or partners involves preparing a reaction mixture containing I5E protein,
20 polypeptide, peptide or fusion protein as described in Sections 5.5.1 and 5.5.2 above, and the binding partner under conditions and for a time sufficient to allow the two to interact and bind, thus forming a complex. In order to test a compound for inhibitory activity, the reaction mixture is
25 prepared in the presence and absence of the test compound. The test compound may be initially included in the reaction mixture, or may be added at a time subsequent to the addition of the I5E moiety and its binding partner. Control reaction mixtures are incubated without the test compound or with a
30 placebo. The formation of any complexes between the I5E moiety and the binding partner is then detected. The formation of a complex in the control reaction, but not in the reaction mixture containing the test compound, indicates that the compound interferes with the interaction of the I5E
35 and the interactive binding partner. Additionally, complex formation within reaction mixtures containing the test

compound and normal I5E protein may also be compared to complex formation within reaction mixtures containing the test compound and a mutant I5E. This comparison may be important in those cases wherein it is desirable to identify 5 compounds that disrupt interactions of mutant but not normal I5Es.

The assay for compounds that interfere with the interaction of the I5E and binding partners can be conducted in a heterogeneous or homogeneous format. Heterogeneous 10 assays involve anchoring either the I5E moiety product or the binding partner onto a solid phase and detecting complexes anchored on the solid phase at the end of the reaction. In homogeneous assays, the entire reaction is carried out in a liquid phase. In either approach, the order of addition of 15 reactants can be varied to obtain different information about the compounds being tested. For example, test compounds that interfere with the interaction by competition can be identified by conducting the reaction in the presence of the test substance; i.e., by adding the test substance to the 20 reaction mixture prior to or simultaneously with the I5E moiety and interactive binding partner. Alternatively, test compounds that disrupt preformed complexes, e.g. compounds with higher binding constants that displace one of the components from the complex, can be tested by adding the test 25 compound to the reaction mixture after complexes have been formed. The various formats are described briefly below.

In a heterogeneous assay system, either the I5E moiety or the interactive binding partner, is anchored onto a solid surface, while the non-anchored species is labeled, either 30 directly or indirectly. In practice, microtitre plates are conveniently utilized. The anchored species may be immobilized by non-covalent or covalent attachments. Non-covalent attachment may be accomplished simply by coating the solid surface with a solution of the I5E gene product or 35 binding partner and drying. Alternatively, an immobilized antibody specific for the species to be anchored may be used

to anchor the species to the solid surface. The surfaces may be prepared in advance and stored.

- In order to conduct the assay, the partner of the immobilized species is exposed to the coated surface with or without the test compound. After the reaction is complete, unreacted components are removed (e.g., by washing) and any complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of ways. Where the non-immobilized species is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the non-immobilized species is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the initially non-immobilized species (the antibody, in turn, may be directly labeled or indirectly labeled with a labeled anti-Ig antibody). Depending upon the order of addition of reaction components, test compounds which inhibit complex formation or which disrupt preformed complexes can be detected.

- Alternatively, the reaction can be conducted in a liquid phase in the presence or absence of the test compound, the reaction products separated from unreacted components, and complexes detected; e.g., using an immobilized antibody specific for one of the binding components to anchor any complexes formed in solution, and a labeled antibody specific for the other partner to detect anchored complexes. Again, depending upon the order of addition of reactants to the liquid phase, test compounds which inhibit complex or which disrupt preformed complexes can be identified.

- In an alternate embodiment of the invention, a homogeneous assay can be used. In this approach, a preformed complex of the I5E moiety and the interactive binding partner is prepared in which either the I5E or its binding partners is labeled, but the signal generated by the label is quenched due to formation of the complex (see, e.g., U.S. Patent No. 4,109,496 by Rubenstein which utilizes this approach for

immunoassays). The addition of a test substance that competes with and displaces one of the species from the preformed complex will result in the generation of a signal above background. In this way, test substances which disrupt I5E/intracellular binding partner interaction can be identified.

In a particular embodiment, an I5E fusion can be prepared for immobilization. For example, the I5E or a peptide fragment, e.g., corresponding to the CD, can be fused to a glutathione-S-transferase (GST) gene using a fusion vector, such as pGEX-5X-1, in such a manner that its binding activity is maintained in the resulting fusion protein. The interactive binding partner can be purified and used to raise a monoclonal antibody, using methods routinely practiced in the art and described above, in Section 5.3. This antibody can be labeled with the radioactive isotope ^{125}I , for example, by methods routinely practiced in the art. In a heterogeneous assay, e.g., the GST-I5E fusion protein can be anchored to glutathione-agarose beads. The interactive binding partner can then be added in the presence or absence of the test compound in a manner that allows interaction and binding to occur. At the end of the reaction period, unbound material can be washed away, and the labeled monoclonal antibody can be added to the system and allowed to bind to the complexed components. The interaction between the I5E gene product and the interactive binding partner can be detected by measuring the amount of radioactivity that remains associated with the glutathione-agarose beads. A successful inhibition of the interaction by the test compound will result in a decrease in measured radioactivity.

Alternatively, the GST-I5E fusion protein and the interactive binding partner can be mixed together in liquid in the absence of the solid glutathione-agarose beads. The test compound can be added either during or after the species are allowed to interact. This mixture can then be added to the glutathione-agarose beads and unbound material is washed away. Again the extent of inhibition of the I5E/binding

partner interaction can be detected by adding the labeled antibody and measuring the radioactivity associated with the beads.

- In another embodiment of the invention, these same techniques can be employed using peptide fragments that correspond to the binding domains of the I5E and/or the interactive or binding partner (in cases where the binding partner is a protein), in place of one or both of the full length proteins. Any number of methods routinely practiced in the art can be used to identify and isolate the binding sites. These methods include, but are not limited to, mutagenesis of the gene encoding one of the proteins and screening for disruption of binding in a co-immunoprecipitation assay. Compensating mutations in the gene encoding the second species in the complex can then be selected. Sequence analysis of the genes encoding the respective proteins will reveal the mutations that correspond to the region of the protein involved in interactive binding. Alternatively, one protein can be anchored to a solid surface using methods described above, and allowed to interact with and bind to its labeled binding partner, which has been treated with a proteolytic enzyme, such as trypsin. After washing, a short, labeled peptide comprising the binding domain may remain associated with the solid material, which can be isolated and identified by amino acid sequencing. Also, once the gene coding for the intracellular binding partner is obtained, short gene segments can be engineered to express peptide fragments of the protein, which can then be tested for binding activity and purified or synthesized.
- For example, and not by way of limitation, an I5E gene product can be anchored to a solid material as described, above, by making a GST-I5E fusion protein and allowing it to bind to glutathione agarose beads. The interactive binding partner can be labeled with a radioactive isotope, such as ³⁵S, and cleaved with a proteolytic enzyme such as trypsin. Cleavage products can then be added to the anchored GST-I5E fusion protein and allowed to bind. After washing away

unbound peptides, labeled bound material, representing the intracellular binding partner binding domain, can be eluted, purified, and analyzed for amino acid sequence by well-known methods. Peptides so identified can be produced 5 synthetically or fused to appropriate facilitative proteins using recombinant DNA technology.

5.4.5. **ASSAYS FOR IDENTIFICATION OF
COMPOUNDS THAT AMELIORATE
I5E RELATED DISORDERS**

10 Compounds, including but not limited to binding compounds identified via assay techniques such as those described, above, in Sections 5.5.1 through 5.5.3, can be tested for the ability to ameliorate I5E disorder symptoms, 15 including inflammatory, central nervous system and metabolic disorders such as body weight disorders. The assays described above can identify compounds which affect I5E activity (e.g., compounds that bind to the I5E, inhibit binding of the natural ligand, and either activate signal transduction (agonists) or block activation (antagonists), 20 and compounds that bind to the natural ligand of the I5E and neutralize ligand activity); or compounds that affect I5E gene activity (by affecting I5E gene expression, including molecules, e.g., proteins or small organic molecules, that affect or interfere with splicing events so that expression 25 of the full length or the truncated form of the I5E can be modulated). However, it should be noted that the assays described can also identify compounds that modulate I5E signal transduction (e.g., compounds which affect downstream signalling events, such as inhibitors or enhancers of G- 30 protein activities which participate in transducing the signal activated by natural I5E ligand binding to the I5E). The identification and use of such compounds which affect another step in the I5E signal transduction pathway in which 35 the I5E gene and/or I5E gene product is involved and, by affecting this same pathway may modulate the effect of I5E on the development of disorders that are within the scope of the

invention. Such compounds can be used as part of a therapeutic method for the treatment of I5E related disorders such as central nervous system inflammatory, immune and metabolic disorders including body weight disorders such as obesity, cachexia and anorexia.

The invention encompasses cell-based and animal model-based assays for the identification of compounds exhibiting such an ability to ameliorate disorder symptoms. Such cell-based assay systems can also be used as the "gold standard" to assay for purity and potency of the natural ligand, natural I5E ligand, including recombinantly or synthetically produced natural I5E ligand and natural I5E ligand mutants.

Cell-based systems can be used to identify compounds which may act to ameliorate gpc-R disorder symptoms. Such cell systems can include, for example, recombinant or non-recombinant cells, such as cell lines, which express the I5E gene. In addition, expression host cells (e.g., COS cells, CHO cells, fibroblasts) genetically engineered to express a functional I5E and to respond to activation by the natural I5E ligand, e.g., as measured by a chemical or phenotypic change, induction of another host cell gene, change in ion flux (e.g., Ca⁺⁺), inositol phosphate levels, tyrosine phosphorylation of host cell proteins, etc., can be used as an end point in the assay.

In utilizing such cell systems, cells may be exposed to a compound suspected of exhibiting an ability to ameliorate I5E disorder symptoms, at a sufficient concentration and for a time sufficient to elicit suppression of disorder symptoms in the exposed cells. After exposure, the cells can be assayed to measure alterations in the expression of the I5E gene, e.g., by assaying cell lysates for I5E mRNA transcripts (e.g., by Northern analysis) or for I5E protein expressed in the cell; compounds which regulate or modulate expression of the I5E gene are good candidates as therapeutics. Still further, the expression and/or activity of components of the signal transduction pathway of which I5E is a part, or the

activity of the I5E signal transduction pathway itself can be assayed.

For example, after exposure, the cell lysates can be assayed for activation of phospholipase C of host cell proteins, as compared to lysates derived from unexposed control cells. The ability of a test compound to inhibit activation of phospholipase C in these assay systems indicates that the test compound inhibits signal transduction initiated by I5E activation. The cell lysates can be readily assayed for phosphatidylinositol turnover as measured by inositol phosphate (IP) accumulation in cells.

In addition, animal-based I5E based disorder systems, which may include, may be used to identify compounds capable of ameliorating disorder-like symptoms. Such animal models may be used as test substrates for the identification of drugs, pharmaceuticals, therapies and interventions which may be effective in treating such disorders. For example, animal models may be exposed to a compound, suspected of exhibiting an ability to ameliorate disorder symptoms, at a sufficient concentration and for a time sufficient to elicit such an amelioration of body symptoms in the exposed animals. The response of the animals to the exposure may be monitored by assessing the reversal of I5E based disorders such as inflammatory, central nervous system and metabolic disorders such as body weight disorders. With regard to intervention, any treatments which reverse any aspect of disorder-like symptoms should be considered as candidates for human therapeutic intervention. Dosages of test agents may be determined by deriving dose-response curves, as discussed in Section 5.7.1, below.

5.5. THE TREATMENT OF I5E BASED DISORDERS

The invention encompasses methods and compositions treating I5E based disorders, including but not limited to immune disorders such as inflammatory disorders, immune and central nervous system disorders. The invention also

encompasses the treatment of metabolic disorders such as body weight disorders, including but not limited to, obesity, cachexia and anorexia.

- Symptoms of certain I5E based disorders may be
5 ameliorated by decreasing the level of I5E gene expression, and/or I5E gene activity, and/or downregulating activity of the I5E pathway (e.g., by targeting downstream signalling events). Different approaches are discussed below.

10 5.5.1. INHIBITION OF I5E EXPRESSION OR I5E ACTIVITY

- Any method which neutralizes natural I5E ligand or inhibits expression of the I5E gene (either transcription or translation) can be used to treat I5E based disorders. Such
15 approaches can be used to treat various maladies such as inflammatory disorders, immune central nervous system disorders or metabolic disorders.

- For example, the administration of soluble peptides, proteins, fusion proteins, or antibodies (including anti-
20 idiotypic antibodies) that bind to and "neutralize" the natural ligand for the I5E, can be used to regulate the I5E. To this end, peptides corresponding to the ECD of I5E, soluble deletion mutants of I5E (e.g., Δ TM I5E mutants), or either of these I5E domains or mutants fused to another
25 polypeptide (e.g., an IgFc polypeptide) can be utilized. Alternatively, anti-idiotypic antibodies or Fab fragments of anti-idiotypic antibodies that mimic the I5E ECD and neutralize natural I5E ligand can be used (see Section 5.3, supra). Such I5E peptides, proteins, fusion proteins, anti-
30 idiotypic antibodies or Fabs are administered to a subject in amounts sufficient to neutralize natural I5E ligand and to inhibit the activity of the I5E.

- Fusion of the I5E, the I5E one or more of the ECDs or the Δ TM I5E to an IgFc polypeptide should not only increase
35 the stability of the preparation, but will increase the half-life and activity of the I5E-Ig fusion protein in vivo. The

Fc region of the Ig portion of the fusion protein may be further modified to reduce immunoglobulin effector function.

- In an alternative embodiment for neutralizing circulating natural I5E ligand, cells that are genetically engineered to express such soluble or secreted forms of I5E may be administered to a patient, whereupon they will serve as "bioreactors" *in vivo* to provide a continuous supply of the natural I5E ligand neutralizing protein. Such cells may be obtained from the patient or an MHC compatible donor and can include, but are not limited to fibroblasts, blood cells (e.g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence for one or more of the I5E ECDs, Δ TM15E, or for I5E-Ig fusion protein (e.g., I5E-, ECD- or Δ TM15E-IgFc fusion proteins) into the cells, e.g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including but not limited to the use of plasmids, cosmids, YACs, electroporation, liposomes, etc. The I5E coding sequence can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve expression and secretion of the I5E peptide or fusion protein. The engineered cells which express and secrete the desired I5E product can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally. Alternatively, the cells can be incorporated into a matrix and implanted in the body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).
- When the cells to be administered are non-autologous cells, they can be administered using well known techniques which prevent the development of a host immune response

against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be
5 recognized by the host immune system.

In an alternate embodiment, therapies can be designed to reduce the level of endogenous I5E gene expression, e.g., using antisense or ribozyme approaches to inhibit or prevent translation of I5E mRNA transcripts; triple helix approaches
10 to inhibit transcription of the I5E gene; or targeted homologous recombination to inactivate or "knock out" the I5E gene or its endogenous promoter. Alternatively, the antisense, ribozyme or DNA constructs described herein could be administered directly to a site containing the target
15 cells; e.g., cells expressing the I5E.

Antisense approaches involve the design of oligonucleotides (either DNA or RNA) that are complementary to I5E mRNA. The antisense oligonucleotides will bind to the complementary I5E mRNA transcripts and prevent translation.
20 Absolutely complementarily, although preferred, is not required. A sequence "complementary" to a portion of an RNA, as referred to herein, means a sequence having sufficient complementarily to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded antisense
25 nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarily and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches
30 with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

35 Oligonucleotides that are complementary to the 5' end of the message, e.g., the 5' untranslated sequence up to and including the AUG initiation codon, should work most

efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have recently shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., 1994, Nature 372:333-335. Thus, oligonucleotides complementary to either the 5'- or 3'- non-translated, non-coding regions of the I5E shown in FIG. 1 could be used in an antisense approach to inhibit translation of endogenous I5E mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5'-, 3'- or coding region of I5E mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

Regardless of the choice of target sequence, it is preferred that *in vitro* studies are first performed to quantitate the ability of the antisense oligonucleotide to inhibit gene expression. It is preferred that these studies utilize controls that distinguish between antisense gene inhibition and nonspecific biological effects of oligonucleotides. It is also preferred that these studies compare levels of the target RNA or protein with that of an internal control RNA or protein. Additionally, it is envisioned that results obtained using the antisense oligonucleotide are compared with those obtained using a control oligonucleotide. It is preferred that the control oligonucleotide is of approximately the same length as the test oligonucleotide and that the nucleotide sequence of the oligonucleotide differs from the antisense sequence no more than is necessary to prevent specific hybridization to the target sequence.

- The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate
- 5 backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors in vivo), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., 1989,
 - 10 Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556; Lemaitre et al., 1987, Proc. Natl. Acad. Sci. 84:648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents.
 - 15 (See, e.g., Krol et al., 1988, BioTechniques 6:958-976) or intercalating agents. (See, e.g., Zon, 1988, Pharm. Res. 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent,
 - 20 hybridization-triggered cleavage agent, etc.

- The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine,
- 25 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine,
 - 30 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine,
 - 35 uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-

5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-0-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330).

Oligonucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

While antisense nucleotides complementary to the I5E coding region sequence could be used, those complementary to the transcribed untranslated region are most preferred. For example, antisense oligonucleotides having the following sequences can be utilized in accordance with the invention:

The antisense molecules should be delivered to cells

which express the I5E *in vivo*. A number of methods have been developed for delivering antisense DNA or RNA to cells; e.g., antisense molecules can be injected directly into the tissue site, or modified antisense molecules, designed to target the 5 desired cells (e.g., antisense linked to peptides or antibodies that specifically bind receptors or antigens expressed on the target cell surface) can be administered systemically.

However, it is often difficult to achieve intracellular concentrations of the antisense sufficient to suppress translation of endogenous mRNAs. Therefore a preferred approach utilizes a recombinant DNA construct in which the antisense oligonucleotide is placed under the control of a strong pol III or pol II promoter. The use of such a construct to transfect target cells in the patient will result in the transcription of sufficient amounts of single stranded RNAs that will form complementary base pairs with the endogenous I5E transcripts and thereby prevent translation of the I5E mRNA. For example, a vector can be introduced in vivo such that it is taken up by a cell and directs the transcription of an antisense RNA. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the antisense RNA can be by any promoter known in the art to act in mammalian, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include but are not limited to: the SV40 early promoter region (Berneist and Chambon, 1981, Nature 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., 1980, Cell 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445), the regulatory sequences of the metallothionein gene

- (Brinster et al., 1982, Nature 296:39-42), etc. Any type of plasmid, cosmid, YAC or viral vector can be used to prepare the recombinant DNA construct which can be introduced directly into the tissue site; e.g., tissue in which I5E is expressed. Alternatively, viral vectors can be used which selectively infect the desired tissue; (e.g., for brain, herpesvirus vectors may be used), in which case administration may be accomplished by another route (e.g., systemically).
- 10 Ribozyme molecules designed to catalytically cleave I5E mRNA transcripts can also be used to prevent translation of I5E mRNA and expression of I5E. (See, e.g., PCT International Publication WO90/11364, published October 4, 1990; Sarver et al., 1990, Science 247:1222-1225). While
- 15 ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy I5E mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole
- 20 requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, 1988, Nature, 334:585-591. There are hundreds of potential
- 25 hammerhead ribozyme cleavage sites within the nucleotide sequence of human I5E cDNA. Preferably the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the I5E mRNA; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional
- 30 mRNA transcripts.
- The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one which occurs naturally in Tetrahymena Thermophila (known as the IVS, or L-19 IVS RNA) and which has been
- 35 extensively described by Thomas Cech and collaborators (Zaug, et al., 1984, Science, 224:574-578; Zaug and Cech, 1986, Science, 231:470-475; Zaug, et al., 1986, Nature, 324:429-

433; published International patent application No. WO 88/04300 by University Patents Inc.; Been and Cech, 1986, Cell, 47:207-216). The Cech-type ribozymes have an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes which target eight base-pair active site sequences that are present in I5E.

- As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express the I5E *in vivo*. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous I5E messages and inhibit translation. Because ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.
- Endogenous I5E gene expression can also be reduced by inactivating or "knocking out" the I5E gene or its promoter using targeted homologous recombination. (E.g., see Smithies et al., 1985, Nature 317:230-234; Thomas & Capecchi, 1987, Cell 51:503-512; Thompson et al., 1989 Cell 5:313-321; each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional I5E (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous I5E gene (either the coding regions or regulatory regions of the I5E gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express I5E *in vivo*. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the I5E gene. Such approaches are particularly suited in the agricultural field where modifications to ES (embryonic stem) cells can be used to generate animal offspring with an inactive I5E (e.g., see Thomas & Capecchi 1987 and Thompson 1989, supra). However

00224303.112800

this approach can be adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site in vivo using appropriate viral vectors, e.g., herpes virus vectors for delivery to brain

5 tissue.

Alternatively, endogenous I5E gene expression can be reduced by targeting deoxyribonucleotide sequences complementary to the regulatory region of the I5E gene (i.e., the I5E promoter and/or enhancers) to form triple helical
10 structures that prevent transcription of the I5E gene in target cells in the body. (See generally, Helene, C. 1991, Anticancer Drug Des., 6(6):569-84; Helene, C., et al., 1992, Ann, N.Y. Accad. Sci., 660:27-36; and Maher, L.J., 1992, Bioassays 14(12):807-15).

15

5.5.2. RESTORATION OR INCREASE IN I5E EXPRESSION OR ACTIVITY

With respect to an increase in the level of normal I5E gene expression and/or I5E gene product activity, I5E nucleic acid sequences can be utilized for the treatment of disorders
20 resulting from a dysfunctional I5E, including inflammatory, central nervous system. Where the cause of a given disorder is a defective I5E, treatment can be administered, for example, in the form of gene replacement therapy.

25 Specifically, one or more copies of a normal I5E gene or a portion of the I5E gene that directs the production of an I5E gene product exhibiting normal function, may be inserted into the appropriate cells within a patient or animal subject, using vectors which include, but are not limited to
30 adenovirus, adeno-associated virus, retrovirus and herpes virus vectors, in addition to other particles that introduce DNA into cells, such as liposomes.

Targeted homologous recombination can be utilized to correct the defective endogenous I5E gene in the appropriate
35 tissue. In animals, targeted homologous recombination can be used to correct the defect in ES cells in order to generate offspring with a corrected trait.

0972392-12000

Additional methods which may be utilized to increase the overall level of I5E gene expression and/or I5E activity include the introduction of appropriate I5E-expressing cells, preferably autologous cells, into a patient at positions and in numbers which are sufficient to ameliorate the symptoms of I5E disorders. Such cells may be either recombinant or non-recombinant. Among the cells which can be administered to increase the overall level of I5E gene expression in a patient are normal cells. The cells can be administered at the anatomical site in the brain, or as part of a tissue graft located at a different site in the body. Such cell-based gene therapy techniques are well known to those skilled in the art, see, e.g., Anderson, et al., U.S. Patent No. 5,399,349; Mulligan & Wilson, U.S. Patent No. 5,460,959.

Finally, compounds, identified in the assays described above, that stimulate or enhance the signal transduced by activated I5E, e.g., by activating downstream signalling proteins in the I5E cascade and thereby by-passing the defective I5E, can be used to treat I5E based disorders. The formulation and mode of administration will depend upon the physico-chemical properties of the compound. The administration should include known techniques that allow for a crossing of the blood-brain barrier.

25 **5.6. PHARMACEUTICAL PREPARATIONS AND METHODS OF ADMINISTRATION**

The compounds that are determined to affect I5E gene expression or I5E activity can be administered to a patient at therapeutically effective doses to treat or ameliorate I5E based disorders including inflammatory, central nervous system and metabolic disorders. A therapeutically effective dose refers to that amount of the compound sufficient to result in amelioration of symptoms of the various disorders.

35

5.6.1. EFFECTIVE DOSE

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD_{50} (the dose lethal to 50% of the population) and the ED_{50} (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD_{50}/ED_{50} . Compounds which exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

15 The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage may

20 vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal

25 models to achieve a circulating plasma concentration range that includes the IC_{50} (i.e., the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in

30 humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

5.6.2. FORMULATIONS AND USE

Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers 5 or excipients.

Thus, the compounds and their physiologically acceptable salts and solvates may be formulated for administration by inhalation or insufflation (either through the mouth or the nose) or oral, buccal, parenteral or rectal administration.

- 10 For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or 15 hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate).
- 20 The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid 25 preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily 30 esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavoring, coloring and sweetening agents as appropriate.
- 35 Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an

acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

10

5.7. DIAGNOSIS DISORDERS ASSOCIATED WITH ABNORMALITIES IN I5E

A variety of methods can be employed for the diagnostic and prognostic evaluation of immune or central nervous system disorders, or metabolic disorders such as body weight disorders, including and for the identification of subjects having a predisposition to such disorders.

Such methods may, for example, utilize reagents such as the I5E nucleotide sequences described in Section 5.1, and I5E antibodies, as described, in Section 5.3. Specifically, such reagents may be used, for example, for: (1) the detection of the presence of I5E gene mutations, or the detection of either over- or under-expression of I5E mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-abundance of I5E gene product relative to the non-disorder state; and (3) the detection of perturbations or abnormalities in the signal transduction pathway mediated by I5E.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one specific I5E nucleotide sequence or I5E antibody reagent described herein, which may be conveniently used, e.g., in clinical settings, to diagnose patients exhibiting disorder abnormalities.

For the detection of I5E mutations, any nucleated cell can be used as a starting source for genomic nucleic acid. For the detection of I5E gene expression or I5E gene

products, any cell type or tissue in which the I5E gene is expressed, may be utilized.

Nucleic acid-based detection techniques are described, below, in Section 5.7.1. Peptide detection techniques are 5 described, below, in Section 5.7.2.

5.7.1. DETECTION OF THE I5E GENE AND TRANSCRIPTS

Mutations within the I5E gene can be detected by
10 utilizing a number of techniques. Nucleic acid from any nucleated cell can be used as the starting point for such assay techniques, and may be isolated according to standard nucleic acid preparation procedures which are well known to those of skill in the art.

DNA may be used in hybridization or amplification assays
15 of biological samples to detect abnormalities involving I5E gene structure, including point mutations, insertions, deletions and chromosomal rearrangements. Such assays may include, but are not limited to, Southern analyses, single
20 stranded conformational polymorphism analyses (SSCP), and PCR analyses.

Such diagnostic methods for the detection of I5E gene-specific mutations can involve for example, contacting and incubating nucleic acids including recombinant DNA molecules,
25 cloned genes or degenerate variants thereof, obtained from a sample, e.g., derived from a patient sample or other appropriate cellular source, with one or more labeled nucleic acid reagents including recombinant DNA molecules, cloned genes or degenerate variants thereof, as described in Section
30 5.1, under conditions favorable for the specific annealing of these reagents to their complementary sequences within the I5E gene. Preferably, the lengths of these nucleic acid reagents are at least 15 to 30 nucleotides. After incubation, all non-annealed nucleic acids are removed from
35 the nucleic acid:I5E molecule hybrid. The presence of nucleic acids which have hybridized, if any such molecules exist, is then detected. Using such a detection scheme, the

007245302 112600

(dA)n-(dG-dT)n short tandem repeats. The average separation of (dC-dA)n-(dG-dT)n blocks is estimated to be 30,000-60,000 bp. Markers which are so closely spaced exhibit a high frequency co-inheritance, and are extremely useful in the identification of genetic mutations, such as, for example, mutations within the I5E gene, and the diagnosis of diseases and disorders related to I5E mutations.

Also, Caskey et al. (U.S. Pat. No. 5,364,759, which is incorporated herein by reference in its entirety) describe a DNA profiling assay for detecting short tri and tetra nucleotide repeat sequences. The process includes extracting the DNA of interest, such as the I5E gene, amplifying the extracted DNA, and labelling the repeat sequences to form a genotypic map of the individual's DNA.

The level of I5E gene expression can also be assayed by detecting and measuring I5E transcription. For example, RNA from a cell type or tissue known, or suspected to express the I5E gene may be isolated and tested utilizing hybridization or PCR techniques such as are described, above. The isolated cells can be derived from cell culture or from a patient. The analysis of cells taken from culture may be a necessary step in the assessment of cells to be used as part of a cell-based gene therapy technique or, alternatively, to test the effect of compounds on the expression of the I5E gene. Such analyses may reveal both quantitative and qualitative aspects of the expression pattern of the I5E gene, including activation or inactivation of I5E gene expression.

In one embodiment of such a detection scheme, cDNAs are synthesized from the RNAs of interest (e.g., by reverse transcription of the RNA molecule into cDNA). A sequence within the cDNA is then used as the template for a nucleic acid amplification reaction, such as a PCR amplification reaction, or the like. The nucleic acid reagents used as synthesis initiation reagents (e.g., primers) in the reverse transcription and nucleic acid amplification steps of this method are chosen from among the I5E nucleic acid reagents described in Section 5.1. The preferred lengths of such

00724002-110000

nucleic acid reagents are at least 9-30 nucleotides. For detection of the amplified product, the nucleic acid amplification may be performed using radioactively or non-radioactively labeled nucleotides. Alternatively, enough amplified product may be made such that the product may be visualized by standard ethidium bromide staining or by utilizing any other suitable nucleic acid staining method.

Additionally, it is possible to perform such I5E gene expression assays "in situ", i.e., directly upon tissue sections (fixed and/or frozen) of patient tissue obtained from biopsies or resections, such that no nucleic acid purification is necessary. Nucleic acid reagents such as those described in Section 5.1 may be used as probes and/or primers for such in situ procedures (See, for example, Nuovo, G.J., 1992, "PCR In Situ Hybridization: Protocols And Applications", Raven Press, NY).

Alternatively, if a sufficient quantity of the appropriate cells can be obtained, standard Northern analysis can be performed to determine the level of mRNA expression of the I5E gene.

5.7.2. DETECTION OF THE I5E GENE PRODUCTS

Antibodies directed against wild type or mutant I5E gene products or conserved variants or peptide fragments thereof, which are discussed, above, in Section 5.3, may also be used as I5E based disorder diagnostics and prognostics, as described herein. Such diagnostic methods, may be used to detect abnormalities in the level of I5E gene expression, or abnormalities in the structure and/or temporal, tissue, cellular, or subcellular location of the I5E, and may be performed in vivo or in vitro, such as, for example, on biopsy tissue.

For example, antibodies directed to epitopes of the I5E ECD can be used in vivo to detect the pattern and level of expression of the I5E in the body. Such antibodies can be labeled, e.g., with a radio-opaque or other appropriate compound and injected into a subject in order to visualize

binding to the I5E expressed in the body using methods such as X-rays, CAT-scans, or MRI. Labeled antibody fragments, e.g., the Fab or single chain antibody comprising the smallest portion of the antigen binding region, are preferred
5 for this purpose to promote crossing the blood-brain barrier.

Additionally, any I5E fusion protein or I5E conjugated protein whose presence can be detected, can be administered. For example, I5E fusion or conjugated proteins labeled with a radio-opaque or other appropriate compound can be
10 administered and visualized in vivo, as discussed, above for labeled antibodies.

Alternatively, immunoassays or fusion protein detection assays, as described above, can be utilized on biopsy and autopsy samples in vitro to permit assessment of the
15 expression pattern of the I5E. Such assays are not confined to the use of antibodies that define one of the I5E ECDs, but can include the use of antibodies directed to epitopes of any of the domains of the I5E, e.g., the ECDs, the TM domains and/or CD. The use of each or all of these labeled
20 antibodies will yield useful information regarding translation and intracellular transport of the I5E to the cell surface, and can identify defects in processing.

The tissue or cell type to be analyzed will generally include those which are known, or suspected, to express the
25 I5E gene. The protein isolation methods employed herein may, for example, be such as those described in Harlow and Lane (Harlow, E. and Lane, D., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York), which is incorporated herein by reference
30 in its entirety. The isolated cells can be derived from cell culture or from a patient. The analysis of cells taken from culture may be a necessary step in the assessment of cells that could be used as part of a cell-based gene therapy technique or, alternatively, to test the effect of compounds
35 on the expression of the I5E gene.

For example, antibodies, or fragments of antibodies, such as those described, above, in Section 5.3, useful in the

present invention may be used to quantitatively or qualitatively detect the presence of I5E gene products or conserved variants or peptide fragments thereof. This can be accomplished, for example, by immunofluorescence techniques
5 employing a fluorescently labeled antibody (see below, this Section) coupled with light microscopic, flow cytometric, or fluorimetric detection. Such techniques are especially preferred if such I5E gene products are expressed on the cell surface.

- 10 The antibodies (or fragments thereof) or natural I5E ligand fusion or conjugated proteins useful in the present invention may, additionally, be employed histologically, as in immunofluorescence, immunoelectron microscopy or non-immuno assays, for in situ detection of I5E gene products or
15 conserved variants or peptide fragments thereof, or for natural I5E ligand binding (in the case of labeled natural I5E ligand fusion protein).

- In situ detection may be accomplished by removing a histological specimen from a patient, and applying thereto a
20 labeled antibody or fusion protein of the present invention. The antibody (or fragment) or fusion protein is preferably applied by overlaying the labeled antibody (or fragment) onto a biological sample. Through the use of such a procedure, it is possible to determine not only the presence of the I5E
25 gene product, or conserved variants or peptide fragments, or natural I5E ligand binding, but also its distribution in the examined tissue. Using the present invention, those of ordinary skill will readily perceive that any of a wide variety of histological methods (such as staining procedures)
30 can be modified in order to achieve such in situ detection.

- Immunoassays and non-immunoassays for I5E gene products or conserved variants or peptide fragments thereof will typically comprise incubating a sample, such as a biological fluid, a tissue extract, freshly harvested cells, or lysates
35 of cells which have been incubated in cell culture, in the presence of a detectably labeled antibody capable of identifying I5E gene products or conserved variants or

peptide fragments thereof, and detecting the bound antibody by any of a number of techniques well-known in the art.

The biological sample may be brought in contact with and immobilized onto a solid phase support or carrier such as 5 nitrocellulose, or other solid support which is capable of immobilizing cells, cell particles or soluble proteins. The support may then be washed with suitable buffers followed by treatment with the detectably labeled I5E antibody or natural I5E ligand fusion protein. The solid phase support may then 10 be washed with the buffer a second time to remove unbound antibody or fusion protein. The amount of bound label on solid support may then be detected by conventional means.

By "solid phase support or carrier" is intended any support capable of binding an antigen or an antibody. Well- 15 known supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the 20 present invention. The support material may have virtually any possible structural configuration so long as the coupled molecule is capable of binding to an antigen or antibody. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test 25 tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Preferred supports include polystyrene beads. Those skilled in the art will know many other suitable carriers for binding antibody or antigen, or will be able to ascertain the same by 30 use of routine experimentation.

The binding activity of a given lot of I5E antibody or natural I5E ligand fusion protein may be determined according to well known methods. Those skilled in the art will be able to determine operative and optimal assay conditions for each 35 determination by employing routine experimentation.

With respect to antibodies, one of the ways in which the I5E antibody can be detectably labeled is by linking the same

03724333-112000

to an enzyme and use in an enzyme immunoassay (EIA) (Voller, A., "The Enzyme Linked Immunosorbent Assay (ELISA)", 1978, Diagnostic Horizons 2:1-7, Microbiological Associates Quarterly Publication, Walkersville, MD); Voller, A. et al., 1978, J. Clin. Pathol. 31:507-520; Butler, J.E., 1981, Meth. Enzymol. 73:482-523; Maggio, E. (ed.), 1980, Enzyme Immunoassay, CRC Press, Boca Raton, FL; Ishikawa, E. et al., (eds.), 1981, Enzyme Immunoassay, Kigaku Shoin, Tokyo). The enzyme which is bound to the antibody will react with an appropriate substrate, preferably a chromogenic substrate, in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorimetric or by visual means. Enzymes which can be used to detectably label the antibody include, but are not limited to, malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate, dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. The detection can be accomplished by colorimetric methods which employ a chromogenic substrate for the enzyme. Detection may also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

Detection may also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the antibodies or antibody fragments, it is possible to detect ISE through the use of a radioimmunoassay (RIA) (see, for example, Weintraub, B., Principles of Radioimmunoassays, Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March, 1986, which is incorporated by reference herein). The radioactive isotope can be detected by such means as the use of a gamma counter or a scintillation counter or by autoradiography.

It is also possible to label the antibody with a fluorescent compound. When the fluorescently labeled

antibody is exposed to light of the proper wave length, its presence can then be detected due to fluorescence. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, 5 phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine.

The antibody can also be detectably labeled using fluorescence emitting metals such as ¹⁵²Eu, or others of the lanthanide series. These metals can be attached to the 10 antibody using such metal chelating groups as diethylenetriaminepentacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

The antibody also can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the 15 chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminol, isoluminol, theromatic acridinium ester, imidazole, 20 acridinium salt and oxalate ester.

Likewise, a bioluminescent compound may be used to label the antibody of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in, which a catalytic protein increases the efficiency of the 25 chemiluminescent reaction. The presence of a bioluminescent protein is determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

30 6. EXAMPLE: ISOLATION AND CHARACTERIZATION OF A NOVEL G-PROTEIN COUPLED RECEPTOR

The following subsection describes the isolation and characterization of a novel human G-protein coupled receptor referred to as I5E. The deduced amino acid sequence of the 35 novel receptor molecule indicates homology with the neuropeptide Y receptor (NPY-2).

6.1. MATERIALS AND METHODS

A sheared BAC library was constructed from murine chromosome 2. The average fragment size was 2kb. Fragments were cloned into the vector pJCP2 for nucleotide sequencing.

- 5 Approximately 800 clones were sequenced with vector primers in order to generate a 4.6 fold sequence coverage of the BAC.

Clones were sequenced by standard automated fluorescent dideoxynucleotide sequencing using dye primer chemistry (Applied Biosystems, Inc., Foster City, CA) on Applied

- 10 Biosystems 373 and 377 sequencers. The DNA sequences were screened to eliminate bacterial, ribosomal and mitochondrial contaminants. Sequence artifacts were also eliminated, such as vector and repetitive element sequences.

- The following primers were used to generate a 877 bp
15 fragment which was used to screen a human fetal brain cDNA library:

5'-TGCTGCTTAAACCTGGGTCGG-3'

5'-GGTGTGTGATTTACTGAGTACCG-3'

- Upon amplification the probe was gel purified and
20 radiolabelled according to standard protocols. Screening was performed on a human fetal brain cDNA library. Hybridization was performed overnight at 50° C. A final washing stringency of 1xSSC/1% SDS at 50° C was achieved. Autoradiography was performed overnight.

- 25 Standard DNA sequencing techniques were utilized for the sequencing and identification of the resulting human I5E gene. The same computer programs as above were used to find identity with the NPY-2 receptor.

30 6.2. RESULTS

- The human I5E sequences were searched against a copy of the GenBank nucleotide database using the BLASTIN program (BLASTIN 1.3MP; Altschul et al., 1990, J. Mol. Biol. 215:403) and a non-redundant protein database with the BLASTX program
35 (BLASTX 1.3MP; Altschul et al., supra). Assembly of overlapping clones into contigs resulted in the identification of one exon which contained the gene of

interest. The gene as shown in FIG. 1, contains an open reading frame of 385 amino acids. The 385 amino acids in the open reading frame were predicted to encode a G-protein coupled receptor using the method of Von Heijne (1990, J.

5 Membrane Biol. 115:195). The protein shows 24% homology with the neuropeptide Y receptor (NPY-2) at the amino acid level. The predicted transmembrane domains span from about amino acids 54-77, 90-112, 140-162, 171-191, 224-244, 274-296 and 312-336.

10

7. EXAMPLE: EXPRESSION OF RECOMBINANT I5E IN COS CELLS

The expression plasmid, I5E HA is derived from the vector pcDNAI/Amp (Invitrogen) and contains the following
15 elements: (i) an SV40 origin of replication; (ii) the ampicillin resistance gene; (iii) the E.coli replication origin; (iv) CMV promoter followed by a polylinker region; and an SV40 intron and polyadenylation site. A DNA fragment encoding the entire human I5E precursor with a HA tag fused
20 in frame at its 3' end is cloned into the polylinker region of the pcDNAI/AMP vector, therefore, placing the expression of the human IE5 protein directly under the control of the CMV promoter. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein as previously
25 described (I. Wilson, H. Niman, R. Heighten, A. Cherenon, M. Connolly, and R. Lerner, 1984, Cell 37:767). The linkage of the HA tag to the I5E protein allows easy detection of the recombinant protein with an antibody that recognizes the HA epitope.

30 The plasmid construction strategy is described as follows:

The DNA sequence encoding for I5E, is constructed by PCR using two primers: containing complementary sequences to an XhoI site, translation stop codon, HA tag and the last 15
35 nucleotides of the I5E coding sequence (not including the stop codon). Therefore, the PCR product contains a HindIII site, I5E coding sequence followed by HA tag fused in frame,

5'-GGGACCCAGAACAGAAACACTA-3'

Upon amplification, the probe was gel purified and radio labeled according to standard protocols.

- Library Construction:** A mouse brain cDNA library was constructed following standard protocols, using RNA isolated as follows: Total RNA was isolated from 300 mouse brain tissue in batches of 100, using the guanidinium isothiocyanate/CsCl method of Chrgwin et al. (1979, Biochemistry 18: 5294) as described by R. Selden in Current Protocols for Molecular Biology (4.2.3 Supplement 14). After quantitation, the RNA was diluted to 1 mg/ml in distilled, deionized water and incubated for 30 min at 37°C with an equal volume of DNase solution (20 mM MgCl₂, 2 mM Dtt, 0.1 units DNase, 0.6 units RNase inhibitor in TE) to remove contaminating DNA. The RNA was extracted with phenol/chloroform/isoamyl, and ethanol precipitated. After quantitation at 260 nm, an aliquot was electrophoresed to check the integrity.

- Poly A+ RNA was isolated using an Oligotex-dT kit (catalog # 70042) from Qiagen (Chatsworth, CA) as described by the manufacturer. After quantitation, the mRNA was ethanol precipitated and resuspended at 1 mg/ml in distilled, deionized, DEPC-treated water.

- cDNA screening:** Screening was performed on the mouse brain libraries described above. Hybridization was performed overnight at 65°C using Rapidhybe buffer manufactured by Gibco BRL according to the manufacturer's protocol. A final washing stringency of 2xSSPE/0.5% SDS 3 times at room temperature, 42°C, and 65°C was used.
- DNA sequences:** Standard DNA sequencing techniques were utilized for the sequencing of the resulting putative murine I5E clones.

8.2. RESULTS

- A mouse genomic sequence, termed skm003b185g09b1, was identified via its homology to G-protein coupled receptor sequences. PCR primers were designed from this clone, and

used to generate a 297 bp probe. The 297 bp probe was used to screen a mouse brain cDNA library, as discussed in Section 8.1, above.

Screening of the mouse cDNA library yielded one independent positive clone, designated famb0333. Sequencing revealed that the clone contained a full-length coding region, which encoded a polypeptide corresponding to a murine I5E homolog. The nucleotide (SEQ. ID. No: 3) and derived amino acid (SEQ. ID. No.: 4) sequences of the murine I5E gene are shown in FIG. 2.

9. EXAMPLE: I5E EXPRESSION ANALYSIS

In the Example presented in this section, the results of a Northern hybridization analysis are described which verify that the I5E gene is expressed in the brain. An in situ hybridization analysis is also described which shows that I5E is expressed in specific areas of the brain.

9.1. MATERIALS AND METHODS

20 Northern analyses: The 297 base pair I5E probe described above was used to probe Northern blots containing total mouse RNA. Northern blots were run on RNA extracted from wild type (C57BL/6J) mice following standard protocols.

25 Tissue Preparation: Brain tissue from wild type mice (C57BL/6J) was removed and frozen with powdered dry ice. Cryostat sections were cut at 10 μ m thickness, mounted on superfrost plus slides, manufactured by VWR, and stored at -80°C.

35 In situ hybridization: Tissue sections were air dried for 20 minutes and then incubated for 10 minutes with 4% PFA/PBS. Sections were then washed with 1x PBS twice for 5 minutes each, incubated with 0.25% acetic anhydride/1 M triethanolamine for 10 minutes, washed again with PBS for 5 minutes, and dehydrated with 70, 80, 95, and 100% ethanol for one minute each, followed by incubation with chloroform for 5 minutes. Hybridization were performed with ³⁵S-radiolabeled

50027-12000

(5×10^7 cpm/ml) cRNA probes encoding a 500 base pair segment of the coding region of the mouse clone ckm300b003h11f1, in the presence of 50% formamide, 10% dextran sulfate, 1x Denhardt's solution, 600 mM NaCl, 10 mM DTT, 0.25% SDS and 100 μ g/ml tRNA for 18 hours at 55°C. After hybridization, slides were washed with 10 mM Tris-HCl (pH 7.6)/500 mM NaCl/1 mM EDTA (TNE) for 10 minutes, incubated in 40 μ g/ml RNase A in TNE at 37°C for 30 minutes, washed in TNE for 10 minutes, incubated once in 2x SSC at 60°C for 1 hour, once in 0.2xSSC at 60°C for 1 hour, in 0.2x SSC at 65°C for 1 hour, and dehydrated with 50, 70, 80, 95, and 100% ethanol. Localization of mRNA transcripts was detected by film emulsion autoradiography, followed by dipping slides in photo-emulsion for precise autoradiographic localization.

15

9.2. RESULTS

Northern analyses were run on RNA extracted from various tissues from wild type (C57BL/6J) mice, using the 297 bp I5E probe described above. Northern analyses showed that murine I5E mRNA transcripts are present in the brain, liver, and spleen.

Subsequently, the 500 base pair segment from ckm300b003h11f1 was used as a probe for an in situ hybridization analysis. Specifically, the antisense cRNA probe was hybridized to sections of wild type mouse (C57BL/6J) brain tissue. mRNA transcripts hybridizing to the antisense cRNA probe were localized to discrete regions of the brain. The results of the in situ hybridization assay are shown in Table I, below. Signal was observed in specific areas of the brain, as shown in Table I, demonstrating that the murine I5E gene is expressed in those tissues.

35

TABLE I

	BRAIN REGIONS	ISE
	Lateral septal n.	+++
5	Septohypothalamic n.	++++
	paraventricular thalamic n. anterior	++++
	superchiasmatic n.	++++
	anterior cortical amygdaloid n.	++
10	piriform cortex	++
	paracentral thalamic n.	+++
	lateral habenular n.	+++
	paraventricular hypothalamic n. (PVN)	++
15	amygdaloid nucleus area	++
	arcuate n.	++
	ventromedial hypothalamic n. (VMH)	+
20		
25		
30		
35		

The following microorganisms were deposited with the American Type Culture Collection (ATCC), Rockville, Maryland on April 18, 1997 and assigned the indicated accession number:

5

<u>Microorganism</u>	<u>ATCC Accession No.</u>
<i>E. Coli</i> , DH10B Ep 065b	98414

10 All publications and patent applications mentioned in the specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

15 The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

25

30

35

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising:
 - a. a nucleic acid molecule encoding a polypeptide comprising the amino acid sequence shown in FIG. 1 (SEQ. ID NO. 2); or
 - b. a nucleic acid molecule encoding a polypeptide comprising the amino acid sequence encoded by the nucleic acid insert of the clone contained in ATCC accession 98414.
2. The isolated nucleic acid molecule of Claim 1 wherein the nucleic acid molecule contains the nucleotide sequence shown in FIG. 1 (SEQ. ID NO. 1).
3. An isolated nucleic acid molecule which hybridizes to the complement of the nucleic acid molecule of Claim 1 and encodes a polypeptide involved in an immune, central nervous system or metabolic disorder.
4. The isolated nucleic acid molecule of Claim 3 wherein the immune disorder is an inflammatory disorder.
5. The isolated nucleic acid molecule of Claim 4 wherein the central nervous system disorder is schizophrenia, cognitive disorders, multiple sclerosis or depression.
6. The isolated nucleic acid molecule of Claim 4 wherein the metabolic disorder is a body weight disorder.
7. An isolated nucleic acid molecule which hybridizes under stringent conditions to the complement of the nucleic acid molecule of Claim 1.
8. The isolated nucleic acid molecule of Claim 3 or 7 wherein the nucleic acid molecule encodes a naturally occurring polypeptide.

9. A nucleotide vector containing the nucleotide sequence of Claim 1, 3 or 7.

10. An expression vector containing the nucleotide sequence of Claim 1, 3 or 7 in operative association with a nucleotide regulatory sequence that controls expression of the nucleotide sequence in a host cell.

11. The expression vector of Claim 10, wherein said regulatory element is selected from the group consisting of the cytomegalovirus hCMV immediate early gene, the early or late promoters of SV40 adenovirus, the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the major operator and promoter regions of phage A, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase, the promoters of acid phosphatase, and the promoters of the yeast α -mating factors.

12. A genetically engineered host cell that contains the nucleotide sequence of Claim 1, 3 or 7.

13. A genetically engineered host cell that contains the nucleotide sequence of Claim 1, 3 or 7 in operative association with a nucleotide regulatory sequence that controls expression of the nucleotide sequence in the host cell.

14. An isolated gene product comprising:
a. the amino acid sequence shown in FIG. 1 (SEQ. ID NO. 2); or
b. the amino acid sequence encoded by the nucleic acid insert of the clone contained in ATCC accession No. 98414.

15. An isolated gene product encoded by the nucleic acid molecule of Claim 3 or 7.

16. An antibody that immunospecifically binds the gene product of Claim 14.

17. An antibody that immunospecifically binds the 5 gene product of Claim 15.

18. A method for diagnosing an immune, central nervous system or metabolic disorder in a mammal, comprising: measuring I5E gene expression in a patient sample.

10

19. The method of Claim 18 wherein the immune disorder is an inflammatory disorder.

20. The method of Claim 18 wherein the central 15 nervous system disorder is schizophrenia, cognitive disorders, multiple sclerosis or depression.

21. The method of Claim 19 wherein the metabolic disorder is a body weight disorder.

20

22. The method of Claim 18 in which expression is measured by detecting I5E mRNA transcripts.

23. The method of Claim 18 in which expression is 25 measured by detecting I5E gene product.

24. A method for diagnosing a I5E disorder in a mammal, comprising: measuring I5E gene expression in a patient sample.

30

25. The method of Claim 24 in which expression is measured by detecting I5E mRNA transcripts.

26. The method of Claim 24 in which expression is 35 measured by detecting I5E gene product.

27. A method for diagnosing an immune, central nervous system or metabolic disorder in a mammal, comprising: detecting a I5E gene mutation contained in the genome of the mammal.

5

28. The method of Claim 27 wherein the central nervous system disorder is schizophrenia, cognitive disorders, multiple sclerosis or depression.

10 29. The method of Claim 27 wherein the immune disorder is an inflammatory disorder.

30. The method of Claim 28 wherein the metabolic disorder is a body weight disorder.

15

31. A method for diagnosing a I5E disorder in a mammal, comprising: detecting a I5E gene mutation contained in the genome of the mammal.

20 32. A method for identifying a compound capable of modulating a I5E activity, comprising:

- a. contacting a compound to a cell that expresses a I5E gene;
- b. measuring the level of I5E gene expression in the cell; and
- 25 c. comparing the level obtained in (b) to I5E gene expression level obtained in the absence of the compound;

such that if the level obtained in (b) differs from that
30 obtained in the absence of the compound, a compound capable of modulating a I5E activity has been identified.

33. The method of Claim 32 wherein the compound increases the level of I5E gene expression.

35

34. The method of Claim 32 wherein the compound decreases the level of I5E gene expression.

35. The method of Claim 32 in which expression of the I5E gene is detected by measuring I5E mRNA transcripts.

36. The method of Claim 32 in which expression of
5 the I5E gene is detected by measuring I5E gene product.

37. The method of Claim 32 wherein the compound is a small organic molecule.

10 38. A method for identifying a compound capable of treating an immune, central nervous system or metabolic disorder, comprising:

- a. contacting a compound to a cell that expresses a I5E gene;
 - 15 b. measuring the level of I5E gene expression in the cell; and
 - c. comparing the level obtained in (b) to I5E gene expression level obtained in the absence of the compound;
- 20 such that if the level obtained in (b) differs from that obtained in the absence of the compound, a compound capable of treating an immune, central nervous system or metabolic disorder has been identified.

25 39. The method of Claim 38 wherein the central nervous system disorder is schizophrenia, cognitive disorders multiple sclerosis or depression.

40. The method of Claim 38 wherein the immune
30 disorder is an inflammatory disorder.

41. The method of Claim 38 wherein the metabolic disorder is a body weight disorder.

35 42. The method of Claim 38 wherein the compound increases the level of I5E gene expression.

43. The method of Claim 38 wherein the compound decreases the level of I5E gene expression.

44. The method of Claim 38 in which expression of the I5E gene is detected by measuring I5E mRNA transcripts.

45. The method of Claim 38 in which expression of the I5E gene is detected by measuring I5E gene product.

46. The method of Claim 38 in which the compound is a small organic molecule.

47. A method for treating an immune, central nervous system or metabolic disorder in a mammal comprising administering to the mammal a compound to the mammal that modulates the synthesis, expression or activity of a mammalian I5E gene or I5E gene product so that symptoms of the disorder are ameliorated.

48. The method of Claim 47 wherein the neuropsychiatric disorder is schizophrenia, attention deficit disorder, a schizoaffective disorder, a bipolar affective disorder or a unipolar disorder.

49. The method of Claim 47 wherein immune disorder is an inflammatory disorder.

50. The method of Claim 47 wherein the metabolic disorder is a body weight disorder.

51. The method of Claim 47 wherein the compound increases the synthesis, expression or activity of a mammalian I5E gene or I5E gene product.

52. The method of Claim 51 wherein the compound comprises the nucleic acid molecule of Claim 1, 3 or 7.

53. The method of Claim 51 wherein the compound is a small organic molecule.

54. The method of Claim 47 wherein the compound decreases the synthesis, expression or activity of a mammalian I5E gene or I5E gene product.

55. The method of Claim 54 wherein the compound provides an antisense or ribozyme molecule that blocks translation of I5E mRNAs.

56. The method of Claim 54 wherein the compound provides a nucleic acid molecule that is complementary to a I5E gene and blocks I5E transcription via triple helix formation.

57. The method of Claim 54 wherein the compound is a small organic molecule.

58. A method for treating a I5E disorder in a mammal comprising administering to the mammal a compound to the mammal that modulates the synthesis, expression or activity of a mammalian I5E gene or I5E gene product so that symptoms of the disorder are ameliorated.

59. The method of Claim 58 wherein the compound increases the synthesis, expression or activity of a mammalian I5E gene or I5E gene product.

60. The method of Claim 59 wherein the compound comprises the nucleic acid molecule of Claim 1, 3 or 7.

61. The method of Claim 59 wherein the compound is a small organic molecule.

62. The method of Claim 58 wherein the compound decreases the synthesis, expression or activity of a mammalian I5E gene or I5E gene product.

5 63. The method of Claim 62 wherein the compound provides an antisense or ribozyme molecule that blocks translation of I5E mRNAs.

64. The method of Claim 62 wherein the compound 10 provides a nucleic acid molecule that is complementary to a I5E gene and blocks I5E transcription via triple helix formation.

65. The method of Claim 62 wherein the compound is 15 a small organic molecule.

66. A method of treating an immune, central nervous system or metabolic disorder resulting from a mutation in a I5E gene, in a mammal, comprising supplying the 20 mammal with a nucleic acid molecule that encodes an unimpaired I5E gene product such that an unimpaired I5E gene product is expressed and symptoms of the disorder are ameliorated.

25 67. The method of Claim 66 wherein the neuropsychiatric disorder is schizophrenia, attention deficit disorder, a schizoaffective disorder, a bipolar affective disorder or a unipolar disorder.

30 68. The method of Claim 66 wherein the immune disorder is an inflammatory disorder.

69. The method of Claim 67 wherein the metabolic disorder is a body weight disorder.

35

70. The method of Claim 66 in which a nucleic acid molecule encoding the unimpaired I5E protein, contained in a

pharmaceutically acceptable carrier, is administered to the mammal.

71. The method of Claim 70 in which the carrier is
5 a DNA vector, a viral vector, a liposome or lipofectin.

72. The method of Claim 66 in which the nucleic acid encoding an unimpaired I5E protein is introduced into the brain of the mammal.

10

73. A method of treating a I5E disorder resulting from a mutation in a I5E gene in a mammal, comprising supplying the mammal with a nucleic acid molecule that encodes an unimpaired I5E gene product such that an
15 unimpaired I5E gene product is expressed and symptoms of the disorder are ameliorated.

74. The method of Claim 73 in which a nucleic acid molecule encoding an unimpaired I5E protein, contained in a
20 pharmaceutically acceptable carrier, is administered to the mammal.

75. The method of Claim 74 in which the carrier is a DNA vector, a viral vector, a liposome or lipofectin.
25

76. A method of treating an immune, central nervous system or metabolic disorder resulting from a mutation in a I5E gene in a mammal, comprising supplying the mammal with a cell comprising a nucleic acid molecule that
30 encodes an unimpaired I5E gene product such that the cell expresses unimpaired I5E gene product and symptoms of the disorder are ameliorated.

77. The method of Claim 76 wherein the
35 neuropsychiatric disorder is schizophrenia, attention deficit disorder, a schizoaffective disorder, a bipolar affective disorder or a unipolar disorder.

78. The method of Claim 76 wherein the immune disorder is inflammation.

5 79. The method of Claim 76 wherein the metabolic disorder is a body weight disorder.

80. The method of Claim 76 in which the cell is engineered ex vivo to express an unimpaired I5E protein.

10

81. The method of Claim 76 in which the cell is contained in a carrier.

82. The method of Claim 76 in which a nucleic acid
15 molecule encoding an unimpaired I5E protein, contained in a pharmaceutically acceptable carrier, is administered to the mammal.

83. The method of Claim 82 in which the carrier is
20 a DNA vector, a viral vector, a liposome or lipofectin.

84. A method of treating a I5E disorder resulting from a mutation in a I5E gene in a mammal, comprising supplying the mammal with a cell comprising a nucleic acid
25 molecule that encodes an unimpaired I5E gene product such that the cell expresses unimpaired I5E gene product and symptoms of the disorder are ameliorated.

85. The method of Claim 84 in which the cell is
30 engineered ex vivo to express an unimpaired I5E protein.

86. The method of Claim 84 in which the cell is contained in a carrier.

35 87. The method of Claim 84 in which a nucleic acid molecule encoding an unimpaired I5E protein, contained in a

pharmaceutically acceptable carrier, is administered to the mammal.

88. The method of Claim 84 in which the carrier is a DNA vector, a viral vector, a liposome or lipofectin.

10

15

20

25

30

35

ABSTRACT

The present invention relates to the discovery, identification and characterization of nucleic acids that
5 encode a novel G protein coupled receptor (I5E) protein. The invention encompasses I5E nucleotides, host cell expression systems, I5E proteins, fusion proteins, polypeptides and peptides, antibodies to the receptor, transgenic animals that express an I5E transgene, or recombinant knock-out animals
10 that do not express the I5E, antagonists and agonists of the receptor, and other compounds that modulate I5E gene expression or I5E activity that can be used for diagnosis, drug screening, clinical trial monitoring, and/or used to treat disorders such as inflammatory, central nervous system
15 or gastrointestinal disorders.

20

25

30

35

CCTGGCCAACTGGCCAAAACCCATCTCTGAGAAAATACAGAAATTAGCTGGGCTGGTGGCACATGCCCTGTGGTCCCAG 3490
CTCCTCAGAGGCGCTGAGGCCATCAGAAATTGCTTGAACCCACAGGGCGAGGGTTCAGTGAATTGAGATGGCACCACTGCA 3569
CTCCAGCGCTGGGTGATAGAGCAGATTCCTCTCAAAAGGAAANTAAAGAAATGCAACACACCTATATATATAGCGCT 3648
AAGCAAACTGTATTATCTGATTTACAAAATCTCTACTTGGCTTGAATGCAATGTGTAAATATGCAATTCAA 3727
AGCCAGCAAGTAACAMTTTTTGGTTATGTACMTTCTGTAAATATAATTAATGTATTTTTATTATATATCTATCTG 3806
GCTCTAGCGGATCAGGAGATCTTTAGGAATGGACCATGTACCAATCAATCTGTCTAGCAGGATCTCATCCCTGTT 3885
CCTTTTGTCTTAGAATATACCACTTCCCTTTCATTTGAAATTTAACTGAAAAACTTTTGTAAATATCAATGTGTATTT 3964
GTGATTTTCCAGTGAATAAAGTGTATGTGTATCCAAATTAATATTTAATCATGTGGATTTAAAAAATAAAAAAAG 4043
GGCGGCCGC 4052

GAATTCCTCCGGGTCGACCCAGCGCTCCGGGCGGGCTGGAACCTCCCGCTTATTGGTCCCGGTGGCGATCTTTGGGAGACCA 79

ATAGACGCCCCAGAGGGAGGACACTGGGATCCAGACTCCACTGGAACCCCGCTTTTCAGATCCTGGATGOTATCTGTTC 158

TCCCTAAGGATTCTAACAGGGACCTGCACCTCACTGACCCAGCAGAAGTGCTGAACCTCCACGTGAGCGCATCTCCCTGA 237

M G P Q N R N T S F A P 12

TACACACCAGCCCACTGTAGCATCATCAAC ATG GGA CCC CAG AAC AGA AAC ACT AGC TTT GCA CCA 304

D L N P P Q D H V S L N Y S Y G D Y D L 32

GAC TTG AAT CCA CCC CAA GAC CAT GTC TCC TTA AAC TAC AGT TAT GGT GAT TAT GAC CTC 364

P L G E D E D V T K T Q T F F A A K I V 52

CCC CTG GGT GAG GAT GAG GAT GTG ACC AAG ACA CAG ACC TTC TTT GCA GCC AAA ATT GTC 424

I G V A L A G I M L V C G I G N F V F I 72

ATT GGC GTG GCA CTG GCA GGC ATC ATG CTG GTC TGC GGC ATT GGC AAC TTT GTC TTC ATT 484

A A L A R Y K K L R N L T N L L I A N L 92

GCT GCC CTC GCC CGC TAC AAG AAG CTG CGC AAC CTT ACC AAC CTC CTC ATT GCT AAC CTG 544

A I S D F L V A I V C C P F F E M D Y Y V 112

GCC ATC TCT GAC TTC CTG GTG GCG ATC GTC TGC TGC CCC TTT GAG ATG GAC TAT TAT GTA 604

V R Q L S W A H G H V L C A S V N Y L R 132

GTA CGG CAG CTT TCC TGG GCG CAT GGT CAC GTG CTT TGT GCC TCC GTC AAC TAC CTT CGT 664

T V S L Y V S T N A L L A I A I D R Y L 152

ACG GTC TCC CTG TAC GTC TCC ACC AAC GCT CTG CTG GCC ATC GCT ATT GAC AGA TAC CTC 724

A I V H P L K P R M N Y Q T A S F L I A 172

GCT ATT GTC CAC CCT TTG AAA CCA CGG ATG AAT TAT CAG ACC GCT TCC TTC CTG ATC GCT 784

L V W M V S I L I A V P S A Y F T T E T 192

TTG GTC TGG ATG GTC TCC ATC CTC ATC GCT GTC CCA TCT GCC TAC TTC ACC ACA GAA ACC 844

I L V I V K N Q E K I F C G Q I W S V D 212

FIG. 2A

05724300-119000

ATC CTC GTT ATC GTC AAG AAT CAA GAA AAA ATC TTC TGT GGT CAG ATC TGG TCG GTG GAC 904

Q Q L Y Y K S Y F L F V F G L E F V G P 232
CAG CAG CTC TAC TAC AAA TCC TAC TTC CTC TTC GTC TTC GGG CTT GAG TTC GTG GGT CCC 944

V V T M T L C Y A R I S Q E L W F K A V 252
GTG GTC ACT ATG ACC CTG TGC TAT GCC AGG ATC TCC CAA GAG CTC TGG TTC AAG GCT GTA 1034

P G F Q T E Q I R K R L R C R R K T V L 272
CCT GGC TTC CAG ACG GAG CAA ATC CGC AAG CGG CTG CGT TGC CGC CGC AAG ACA GTG CTA 1044

L L M G I L T A Y V L C W A P F Y G F T 292
CTG CTC ATG GGC ATC CTC ACA GCC TAC GTG CTG TGC TGG GCG CCG TTC TAT GGC TTT ACC 1144

I V R D F F P T V V V K E K H Y L T A F 312
ATA GTG CGA GAC TTC TTC CCC ACG GTA GTT GTG AAG GAG AAG CAC TAC CTC ACC GCC TTC 1204

Y V V E C I A M S N S M I N T I C F V T 332
TAC GTC GTG GAG TGC ATT GCC ATG AGC AAC AGC ATG ATC AAT ACT ATA TGC TTC GTG ACG 1264

V K N N T M K Y F K K M L R L H W R P S 352
GTC AAG AAC AAC ACC ATG AAA TAC TTC AAG AAG ATG CTG CGG CTC CAC TGG CGG CCC TCT 1324

H Y G S K S S A D L D L K T S G V P A T 372
CAC TAC GGG AGT AAG TCC AGC GCT GAC CTC GAC CTC AAA ACC AGC GGG GTG CCT GCC ACT 1384

E E V D C I R L K * 381
GAA GAG GTG GAT TGT ATC AGA CTA AAG TAG 1444

CCTTCAGGTGTGCCCAAGGAAAAATTAACATTCCGGTACTCAGTAAATCACACACCATCAACCACTCAAGCTACAT 1493

GGAAAGATACGGCTGTATTACGTTCTCCTGCTCTAATGTATCAGGACGCTTCTATGTAATAACATACAGCACAACTGA 1572

TGTCTGCATAACATCTTTAGAGGCAGACACAAATAGTAACAAGTGATGTGGACTGAATGCTTCTGTCTGCACAAACACAC 1651

CAACCAATTATTCAAGGACAAGAGCTGACATGTGAGAATTACCTGCTATGTGCAAAAACAAGTTACCCCCCAAAAAAT 1730

GATAGAAGCTATTTGGAGTTATTACGCTCTATCTATCTATCTATCTATCCATCCATCCATCCATCCAGGCTACTA 1809

GAAAGAAGTCACAAATGACTAGCCGAGTCACTACATATTCTTTCATTCTGTATCTTTTCTGCACAGAACTGTCAA 1888

Fig. 2B

GGCAATAGATAAAGCACCTAGACATAC'TAGAAATGTAAGGATAACTCCATCAATAGGGAGACCAAGGCCTCATAGGAA 197

GAGGGTCCATATAGTATACTGACTTTCCCCACTCCACACCAGTTATCTCCTTAGATATTCTGTACTTATCTGCAATGTT 2046

GTAATTTCAAATGAGGAAAAATAAGGGGACAGGCTTTACCACAGATGTATCAAATCTCATCAAGCCCATAGGGGCAAGAA 2145

TGGGAGGCTCTGACACAAAGAAATGTATCCAGTTCTGGATAACTTTAATGCCAAGCATTTTCAGGGCTCTGGGGTCTTGG 2204

AGGAAGAGGACACAGAAAGAGCCGAGGTTCCAGTGGCAATGAGTATAATCTGTCCATTTGCTATGATTTGGACAATTT 2283

TCTAGAACATACTCCGACTTACAAAAGGAACCTACTTGTAGATCCAAAGATCCGGGTAAAGTCCTAACCCAGGACTC 2342

ATCTCTGTGTGTCTCCACTGTGAATGAAATGGAATAATGAAAACGGATCATTTAGGAACATCAGCCCGCGAAGTCATGG 2411

TGTGGATGTGATTTTACCTCTTCTCTTGTGAAGAATGAGGTCGTGAAAAGCTCATTTAGAGGGAGTTTGGAAATGGAGAA 2520

ACAGTCCACACTTTTCATCCCTCTTCTTTGAATCGGAGACCACTAAACGCATCTTTGAAGTAGCGTATCTATATGAGG 2599

GCATAAAGGTCTCCCTGTACAGAGTGCATCAAGAAAATACAGTCAATGCCATACCCTCAGCATCCCTGTGGTCTTA 2678

GACAGTCTTCCCAACAAAGCACTGTGTGGACCCCGAGGACTGAATTCACTTGTATTATGTCTATCTACTGAATACTAGG 2757

GTTGATCAAGTTGGCTAGATAGGTATTTCTTCTCCTCTCACAACCCCATATGTATCCCTCCCTTAAATCCAGTTACTA 2836

AGGAAGACCTTCTTAAACACAGGAGAACCATTAATTCTGTCCAGGACACAAATAACCTCTCCAGTAGACACTGTACCCTT 2915

CACATGTCAACAGAAATTGCCTCCTTCTGTATTAAACATATCATCCTCCTTTCATTTAGATTTAACAGAAACCATTT 2994

CCTGTAAATTTCAATGTGTTTGTGATACCGCACTGTAAAAAGCGTATGCTGTTATCATATGGAATAATTAACATACAGA 3073

ATTGTAATCGTAGTTCCTCAAAGGTTCCCTACTCCTGTGTATCTTATGTTTATATGTTTGTATGTAATGGAGCTGTGT 3152

AGCTGTCTAAGCAGCTCAAGCTGAAATGAGGGAATGTCCAATGGTGTCTTGTAGAGCAGGGCCATCTCAGGCTAGCAGC 3231

TGGCCTCAGTCTGTCTCTCTCGGGAGTGTGTTCTTAAATATGAATTAGCAGCAAAACCATTAAAAAAGGG 3300

CGGCCGC 3347

Fig. 2C

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below at 201 et seq. underneath my name.

I believe I am the original, first and sole inventor if only one name is listed at 201 below, or an original, first and joint inventor if plural names are listed at 201 et seq. below, of the subject matter which is claimed and for which a patent is sought on the invention entitled

A NOVEL HUMAN G-PROTEIN COUPLED RECEPTOR

and for which a patent application:

- ☐ is attached hereto and includes amendment(s) filed on (if applicable)
- ☐ was filed in the United States on as Application No. (for declaration not accompanying application) with amendment(s) filed on (if applicable)
- ☐ was filed as PCT international Application No. on and was amended under PCT Article 19 on (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified application, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

EARLIEST FOREIGN APPLICATION(S), IF ANY, FILED PRIOR TO THE FILING DATE OF THE APPLICATION			
APPLICATION NUMBER	COUNTRY	DATE OF FILING (day, month, year)	PRIORITY CLAIMED
			YES <input type="checkbox"/> NO <input type="checkbox"/>
			YES <input type="checkbox"/> NO <input type="checkbox"/>

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

APPLICATION NUMBER	FILING DATE

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NO.	FILING DATE	STATUS		
		PATENTED	PENDING	ABANDONED
08/833,226	April 17, 1997	X		
09/062,753	April 17, 1998		X	

POWER OF ATTORNEY: As a named inventor, I hereby appoint S. Leshe Misrock (Reg. No. 18872), Berj A. Terzian (Reg. No. 20060), David Weid, III (Reg. No. 21094), Jonathan A. Marshall (Reg. No. 24614), Barry D. Rein (Reg. No. 22411), Stanton T. Lawrence, III (Reg. No. 25736), Charles E. McKenney (Reg. No. 22795), Philip T. Shannon (Reg. No. 24278), Francis E. Morris (Reg. No. 24615), Charles E. Miller (Reg. No. 24576), Gidon D. Stern (Reg. No. 27469), John J. Lauter, Jr. (Reg. No. 27814), Brian M. Possant (Reg. No. 28452), Brian D. Coggio (Reg. No. 27624), Rory J. Radding (Reg. No. 28749), Stephen J. Harbulak (Reg. No. 29166), Donald J. Goodell (Reg. No. 19766), James N. Palik (Reg. No. 25510), Thomas E. Friebe (Reg. No. 29258), Laura A. Coruzzi (Reg. No. 30742), Jennifer Gordon (Reg. No. 30753), Allan A. Fanucci (Reg. No. 30256), Geraldine F. Baldwin (Reg. No. 31232), Victor N. Balancia (Reg. No. 31231), Samuel B. Abrams (Reg. No. 30605), Steven I. Wallach (Reg. No. 35402), Marcia H. Sundeen (Reg. No. 30893), Paul J. Zegger (Reg. No. 33821), Edmond R. Bannan (Reg. No. 32110), Bruce J. Barker (Reg. No. 33291), Adriane M. Antler (Reg. No. 32605), Thomas G. Rowan (Reg. No. 34419), James G. Markey (Reg. No. 31636), Thomas D. Kohler (Reg. No. 32797), Scott D. Stimpson (Reg. No. 33607), Gary S. Williams (Reg. No. 31066), William S. Gailiani (Reg. No. 33885), Ann L. Gnsoli (Reg. No. 31956), Todd A. Wagner (Reg. No. 35399), Scott B. Familant (Reg. No. 35514), Kelly D. Talcott (Reg. No. 39582), Francis D. Cerreto (Reg. No. 38100), Anthony M. Insigna (Reg. No. 35203), Brian M. Rothery (Reg. No. 35340), Brian D. Siff (Reg. No. 35679), and Alan Tenenbaum (Reg. No. 34939), all of Pennie & Edmonds LLP, whose addresses are 1155 Avenue of the Americas, New York, New York 10036, 1667 K Street N.W., Washington, DC 20006 and 3300 Hillview Avenue, Palo Alto, CA 94304, and each of them, my attorneys, to prosecute this application, and to transact all business in the Patent and Trademark Office connected therewith.

SEND CORRESPONDENCE TO:		PENNIE & EDMONDS LLP 1155 Avenue of the Americas New York, N.Y. 10036-2711		DIRECT TELEPHONE CALLS TO: PENNIE & EDMONDS LLP DOCKETING (212) 790-2803			
2 0 1	FULL NAME OF INVENTOR	LAST NAME	MOORE	FIRST NAME	KAREN	MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY	MAYNARD	STATE OR FOREIGN COUNTRY	MASSACHUSETTS	COUNTRY OF CITIZENSHIP GREAT BRITAIN	
	POST OFFICE ADDRESS	STREET	34 CHANDLER STREET	CITY	MAYNARD	STATE OR COUNTRY	MA ZIP CODE 01754
2 0 2	FULL NAME OF INVENTOR	LAST NAME	NAGLE	FIRST NAME	DEBORAH	MIDDLE NAME	LYNN
	RESIDENCE & CITIZENSHIP	CITY	WATERTOWN	STATE OR FOREIGN COUNTRY	MASSACHUSETTS	COUNTRY OF CITIZENSHIP UNITED STATES	
	POST OFFICE ADDRESS	STREET	370 ARLINGTON STREET	CITY	WATERTOWN	STATE OR COUNTRY	MA ZIP CODE 02172
2 0 3	FULL NAME OF INVENTOR	LAST NAME	WOOLF	FIRST NAME	ELIZABETH	MIDDLE NAME	A.
	RESIDENCE & CITIZENSHIP	CITY	GEORGETOWN	STATE OR FOREIGN COUNTRY	MASSACHUSETTS	COUNTRY OF CITIZENSHIP UNITED STATES	
	POST OFFICE ADDRESS	STREET	9 BALDPATE ROAD	CITY	GEORGETOWN	STATE OR COUNTRY	MA DHS/SP CODE 01833
2 0 4	FULL NAME OF INVENTOR	LAST NAME		FIRST NAME		MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY		STATE OR FOREIGN COUNTRY		COUNTRY OF CITIZENSHIP	
	POST OFFICE ADDRESS	STREET		CITY		STATE OR COUNTRY	ZIP CODE
2 0 5	FULL NAME OF INVENTOR	LAST NAME		FIRST NAME		MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY		STATE OR FOREIGN COUNTRY		COUNTRY OF CITIZENSHIP	
	POST OFFICE ADDRESS	STREET		CITY		STATE OR COUNTRY	ZIP CODE
2 0 6	FULL NAME OF INVENTOR	LAST NAME		FIRST NAME		MIDDLE NAME	
	RESIDENCE & CITIZENSHIP	CITY		STATE OR FOREIGN COUNTRY		COUNTRY OF CITIZENSHIP	
	POST OFFICE ADDRESS	STREET		CITY		STATE OR COUNTRY	ZIP CODE

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

SIGNATURE OF INVENTOR 201	SIGNATURE OF INVENTOR 202	SIGNATURE OF INVENTOR 203
Karen Moore	Deborah Lynn Nagle	Elizabeth A. Woolf
DATE	DATE	DATE
SIGNATURE OF INVENTOR 204	SIGNATURE OF INVENTOR 205	SIGNATURE OF INVENTOR 206
DATE	DATE	DATE

SEQUENCE LISTING

<110> Moore, K.
Nagle, D.
Woolf, E.

<120> NOVEL HUMAN G-PROTEIN COUPLED RECEPTOR

<130> 7853-119

<140> 09/062,753

<141> 1998-04-17

<150> 08/833,226

<151> 1997-04-17

<160> 8

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 4052

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (45)...(1196)

<400> 1

agccgcagag cgcacagaaa ggaggcgccg agacagacat cacc atg gca gcc cag	56
Met Ala Ala Gln	
1	
aat gga aac acc agt ttc aca ccc aac ttt aat cca ccc caa gac cat	104
Asn Gly Asn Thr Ser Phe Thr Pro Asn Phe Asn Pro Pro Gln Asp His	
5 10 15 20	
gcc tcc tcc ctc tcc ttt aac ttc agt tat ggt gat tat gac ctc cct	152
Ala Ser Ser Leu Ser Phe Asn Phe Ser Tyr Gly Asp Tyr Asp Leu Pro	
25 30 35	
atg gat gag gat gag gac atg acc aag acc cgg acc ttc ttc gca gcc	200
Met Asp Glu Asp Glu Asp Met Thr Lys Thr Arg Thr Phe Phe Ala Ala	
40 45 50	
aag atc gtc att ggc att gca ctg gca ggc atc atg ctg gtc tgc ggc	248
Lys Ile Val Ile Gly Ile Ala Leu Ala Gly Ile Met Leu Val Cys Gly	
55 60 65	
atc ggt aac ttt gtc ttt atc gct gcc ctc acc cgc tat aag aag ttg	296
Ile Gly Asn Phe Val Phe Ile Ala Ala Leu Thr Arg Tyr Lys Lys Leu	
70 75 80	

cgc aac ctc acc aat ctg ctc att gcc aac ctg gcc atc tcc gac ttc 344
 Arg Asn Leu Thr Asn Leu Leu Ile Ala Asn Leu Ala Ile Ser Asp Phe
 85 90 95 100

 ctg gtg gcc atc atc tgc tgc ccc ttc gag atg gac tac tac gtg gta 392
 Leu Val Ala Ile Ile Cys Cys Pro Phe Glu Met Asp Tyr Tyr Val Val
 105 110 115

 cgg cag ctc tcc tgg gag cat ggc cac gtg ctc tgt gcc tcc gtc aac 440
 Arg Gln Leu Ser Trp Glu His Gly His Val Leu Cys Ala Ser Val Asn
 120 125 130

 tac ctg cgc acc gtc tcc ctc tac gtc tcc acc aat gcc ttg ctg gcc 488
 Tyr Leu Arg Thr Val Ser Leu Tyr Val Ser Thr Asn Ala Leu Leu Ala
 135 140 145

 att gcc att gac aga tat ctc gcc atc gtt cac ccc ttg aaa cca cgg 536
 Ile Ala Ile Asp Arg Tyr Leu Ala Ile Val His Pro Leu Lys Pro Arg
 150 155 160

 atg aat tat caa acg gcc tcc ttc ctg atc gcc ttg gtc tgg atg gtg 584
 Met Asn Tyr Gln Thr Ala Ser Phe Leu Ile Ala Leu Val Trp Met Val
 165 170 175 180

 tcc att ctc att gcc atc cca tgc gct tac ttt gca aca gaa acc gtc 632
 Ser Ile Leu Ile Ala Ile Pro Ser Ala Tyr Phe Ala Thr Glu Thr Val
 185 190 195

 ctc ttt att gtc aag agc cag gag aag atc ttc tgt ggc cag atc tgg 680
 Leu Phe Ile Val Lys Ser Gln Glu Lys Ile Phe Cys Gly Gln Ile Trp
 200 205 210

 cct gtg gat cag cag ctc tac tac aag tcc tac ttc ctc ttc atc ttt 728
 Pro Val Asp Gln Gln Leu Tyr Tyr Lys Ser Tyr Phe Leu Phe Ile Phe
 215 220 225

 ggt gtc gag ttc gtg ggc cct gtg gtc acc atg acc ctg tgc tat gcc 776
 Gly Val Glu Phe Val Gly Pro Val Val Thr Met Thr Leu Cys Tyr Ala
 230 235 240

 agg atc tcc cgg gag ctc tgg ttc aag gca gtc cct ggg ttc cag acg 824
 Arg Ile Ser Arg Glu Leu Trp Phe Lys Ala Val Pro Gly Phe Gln Thr
 245 250 255 260

 gag cag att cgc aag cgg ctg cgc tgc cgc agg aag acg gtc ctg gtg 872
 Glu Gln Ile Arg Lys Arg Leu Arg Cys Arg Arg Lys Thr Val Leu Val
 265 270 275

 ctc atg tgc att ctc acg gcc tat gtg ctg tgc tgg gca ccc ttc tac 920
 Leu Met Cys Ile Leu Thr Ala Tyr Val Leu Cys Trp Ala Pro Phe Tyr
 280 285 290

 ggt ttc acc atc gtt cgt gac ttc ttc ccc act gtg ttc gtg aag gaa 968
 Gly Phe Thr Ile Val Arg Asp Phe Phe Pro Thr Val Phe Val Lys Glu
 295 300 305

aag cac tac ctc act gcc ttc tac gtg gtc gag tgc atc gcc atg agc Lys His Tyr Leu Thr Ala Phe Tyr Val Val Glu Cys Ile Ala Met Ser	1016
310 315 320	
aac agc atg atc aac acc gtg tgc ttc gtg acg gtc aag aac aac acc Asn Ser Met Ile Asn Thr Val Cys Phe Val Thr Val Lys Asn Asn Thr	1064
325 330 335 340	
atg aag tac ttc aag aag atg atg ctg ctg cac tgg cgt ccc tcc cag Met Lys Tyr Phe Lys Lys Met Met Leu Leu His Trp Arg Pro Ser Gln	1112
345 350 355	
cgg ggg agc aag tcc agt gct gac ctt gac ctc aga acc aac ggg gtg Arg Gly Ser Lys Ser Ser Ala Asp Leu Asp Leu Arg Thr Asn Gly Val	1160
360 365 370	
ccc acc aca gaa gaa gtg gac tgt atc agg ctg aag tgaccacctg Pro Thr Thr Glu Glu Val Asp Cys Ile Arg Leu Lys	1206
375 380	
gtgtcacaca attgaaaacc ccagtcacgt actcagagca taccaccacca tcaaccaagt tcataggtctg catgggaatat gacatctgtg ttcagtgcct ccccgctgcc tcaagaagcc gaagtctgca aagctcgtaac atacaatgag actagacatg aaccaaaatca gtcgacattt actgatatcc gctcgacacc tactgtgtcc acaatcccca caaggagatt agcacacaagg agcagcaact gacatggact gaacatgtac tgtgtgcaaa ccacaccaat gagattagac ggggacagca gaggctgaca ttactcttc acctactgta atcaaaaaca cttgatttga ttacaatcaa aaacatataa aaacataaac aaagttagcag aagctattgg agtttccaag ctatctccag atatatagat agttcacctt ccactctccc taattctgta tcttaccagt cgaggaatat caaaaaggcta tagggccaggc atgatggctc atgcctgtaa tcccagcact tggggaggct gaggcagctg gatcacttga ggtagagggt tcaaccaggc ctggccaaca tgggtgaaacc ctgtctctac taaaataaca aaattagcta ggcgtgtgtg cgggcgctgt taatcccgat tactcaggag gctgaagcag gagaatagct tgaacctggg agttggagtt tgcatgtgac tgagattgct ccactgcact ccagcctgag tgacagagtg agactctgtc tcaggaaaaa acaacaacaa caaacaacaa acaacaacaa caaacaacaa caaacaacgg ctatagaaga agactcttcc acacaatgga aatgtaacga taagtgtgtc agtgcgtgtg ttacagcata atggggaggtg cgttacagcc atcactactga actttcccac ccactctcta ctgcctccca gggcattctc taggattttg gcttcaagaa aaaaaaaatt cttatagcta gcccagcctt atgtgggtta ccacaatggt gtaatttcaa aggaagaac ctaaaaatca ctttccact gatgcttgaa agcttatcat tttatttggg ttggagatggg taatcctgag gtgtcaattt ttgcctcttc agtgcaaaagg atttcagttg ctctggggct agggggaaag aggacagaga aaaaagtgga ggttgcact ggcaatgga ataatctctg ttgagcattt gctaaggact ggaccacttt ctgaacact cctctcttta caaaaggaac tctacctaga atccaaagac ctgggttcag gtctaaact taagactcaa gtctcaaat catgatgttt tctctgtgtg tctcagtttt gccttaatga aatggcgatg atgaaaatat ctgctctcca taccttgcaa gactgttggg agagcccat gaggccatgg tttgtgaatg tgttttcaa ctgtgcacac gataagaatg gagaagtgat attgaacagt ttatttggag ggagtttatt tggaaacccc atccactgtg atttattaga gaaataccca cacttttcca tccctgtctc ttggatgaaa gactcctgaa gacttcacag tgcacttgg ctacagtgtg ccaaaaaggg atccctgttc ttggttataa tctgggaaat ttaacctcag atctctcagt acccaagac tctcagatc cctcggtctg tagaagtgtt gacagctctc ctgcactgtt gcctaatagc accctagtg tgcataaata tcaactctga atctgtttgt attattatac atttgttgta actgtaggt cactgtctta tttctctctg attcattttg atgtgtgtag tatgaaaagt tgactcgtgt tggagctgac ccatccatat ttgaccaatt ctaatttttt tatgacaag gaattaatgt tttgtctgtg tgattgtttc tattatttgt tgatttgttt cctgtactga agtttcaacc aatgtttctt tctatcacca cccagcagat tcactctcag ccaactcat gtactctcag aaaaatcgag ccggcatggt ggctcacat tgtaatccca gcaactcggg aggccaagat gggcagatac cctgaggtca ggagttcaag accagctcag ccaactggc aaaaacccat ctctagaaaa atacagaaat tagctggcgt ggtggcacat gcctgtgtgc	1266 1326 1386 1446 1506 1566 1626 1686 1746 1806 1866 1926 1986 2046 2106 2166 2226 2286 2346 2406 2466 2526 2586 2646 2706 2766 2826 2886 2946 3006 3066 3126 3186 3246 3306 3366 3426 3486

0974392-112830

ccagctcttc agggagctga ggcattgagaa ttgcttgaac cccagaggca gaggttgacg 3546
tgaattgaga tgcgaccact gcactccagc ctgggtgata gagcaagatt ccattctcaa 3606
agggaaaataa aagaaaatgc aaacacacta taattattagc ctaagcaaaa ctgttaattc 3666
tgattttacaa aaattcttac ttgcttggct ttgaaatgca ttgtgttaata atgcatttca 3726
aagccaagca agtaacaatt ttagggtatg tacatttcta taaataaat aattgtattt 3786
ttattttatta ttctatcttg gctcttagcc gaatcaggag attctttagg aatggaccat 3846
gtaccagtagc agtctgtcag caggattcat caccctgttc ctttttgtcc tagaataatac 3906
caacttctctt tcattgaaat ttaactgaaa aaacttttgt aaataatcagt gtgtatttgt 3966
gtattttccag tgattaaagt gtgactgttgt tatccaatta aataatatac atgtggaatt 4026
taaaaaaaaaa aaaaaagggc ggccgc 4052

<210> 2
<211> 384
<212> PRT
<213> Homo sapiens

<400> 2
Met Ala Ala Gln Asn Gly Asn Thr Ser Phe Thr Pro Asn Phe Asn Pro
1 5 10 15
Pro Gln Asp His Ala Ser Ser Leu Ser Phe Asn Phe Ser Tyr Gly Asp
20 25 30
Tyr Asp Leu Pro Met Asp Glu Asp Glu Asp Met Thr Lys Thr Arg Thr
35 40 45
Phe Phe Ala Ala Lys Ile Val Ile Gly Ile Ala Leu Ala Gly Ile Met
50 55 60
Leu Val Cys Gly Ile Gly Asn Phe Val Phe Ile Ala Ala Leu Thr Arg
65 70 75 80
Tyr Lys Lys Leu Arg Asn Leu Thr Asn Leu Leu Ile Ala Asn Leu Ala
85 90 95
Ile Ser Asp Phe Leu Val Ala Ile Ile Cys Cys Pro Phe Glu Met Asp
100 105 110
Tyr Tyr Val Val Arg Gln Leu Ser Trp Glu His Gly His Val Leu Cys
115 120 125
Ala Ser Val Asn Tyr Leu Arg Thr Val Ser Leu Tyr Val Ser Thr Asn
130 135 140
Ala Leu Leu Ala Ile Ala Ile Asp Arg Tyr Leu Ala Ile Val His Pro
145 150 155 160
Leu Lys Pro Arg Met Asn Tyr Gln Thr Ala Ser Phe Leu Ile Ala Leu
165 170 175
Val Trp Met Val Ser Ile Leu Ile Ala Ile Pro Ser Ala Tyr Phe Ala
180 185 190
Thr Glu Thr Val Leu Phe Ile Val Lys Ser Gln Glu Lys Ile Phe Cys
195 200 205
Gly Gln Ile Trp Pro Val Asp Gln Gln Leu Tyr Tyr Lys Ser Tyr Phe
210 215 220
Leu Phe Ile Phe Gly Val Glu Phe Val Gly Pro Val Val Thr Met Thr
225 230 235 240
Leu Cys Tyr Ala Arg Ile Ser Arg Glu Leu Trp Phe Lys Ala Val Pro
245 250 255
Gly Phe Gln Thr Glu Gln Ile Arg Lys Arg Leu Arg Cys Arg Arg Lys
260 265 270
Thr Val Leu Val Leu Met Cys Ile Leu Thr Ala Tyr Val Leu Cys Trp
275 280 285
Ala Pro Phe Tyr Gly Phe Thr Ile Val Arg Asp Phe Phe Pro Thr Val
290 295 300
Phe Val Lys Glu Lys His Tyr Leu Thr Ala Phe Tyr Val Val Glu Cys
305 310 315 320

Ile Ala Met Ser Asn Ser Met Ile Asn Thr Val Cys Phe Val Thr Val
 325 330 335
 Lys Asn Asn Thr Met Lys Tyr Phe Lys Lys Met Met Leu Leu His Trp
 340 345 350
 Arg Pro Ser Gln Arg Gly Ser Lys Ser Ser Ala Asp Leu Asp Leu Arg
 355 360 365
 Thr Asn Gly Val Pro Thr Thr Glu Glu Val Asp Cys Ile Arg Leu Lys
 370 375 380

<210> 3
 <211> 3317
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (269)...(1411)

<400> 3
 gaattcccggt gtcgaccac ggcgccgggc ggctggaact cccgcttatt ggcccccggt 60
 ggcgatcttt gggagaccaa tagacgcccc agaggaggga cactgggata cagactccac 120
 tggaaaccccg cttttcagat cctggatggt atctgttctc cctaaggatt ctaacaggga 180
 cctgcactca ctgaccccgag cagaagtgtc gaactccacg tgagcgcata tcctgatac 240
 acaccagccc acctgtagca tcataaac atg gga ccc cag aac aga aac act 292
 Met Gly Pro Gln Asn Arg Asn Thr
 1 5
 agc ttt gca cca gac ttg aat cca ccc caa gac cat gtc tcc tta aac 340
 Ser Phe Ala Pro Asp Leu Asn Pro Pro Gln Asp His Val Ser Leu Asn
 10 15 20
 tac agt tat ggt gat tat gac ctc ccc ctg ggt gag gat gag gat gtg 388
 Tyr Ser Tyr Gly Asp Tyr Asp Leu Pro Leu Gly Glu Asp Glu Asp Val
 25 30 35 40
 acc aag aca cag acc ttc ttt gca gcc aaa att gtc att ggc gtg gca 436
 Thr Lys Thr Gln Thr Phe Phe Ala Ala Lys Ile Val Ile Gly Val Ala
 45 50 55
 ctg gca ggc atc atg ctg gtc tgc ggc att ggc aac ttt gtc ttc att 484
 Leu Ala Gly Ile Met Leu Val Cys Gly Ile Gly Asn Phe Val Phe Ile
 60 65 70
 gct gcc ctc gcc cgc tac aag aag ctg cgc aac ett acc aac ctc ctc 532
 Ala Ala Leu Ala Arg Tyr Lys Lys Leu Arg Asn Leu Thr Asn Leu Leu
 75 80 85
 att gct aac ctg gcc atc tct gac ttc ctg gtg gcg atc gtc tgc tgc 580
 Ile Ala Asn Leu Ala Ile Ser Asp Phe Leu Val Ala Ile Val Cys Cys
 90 95 100
 ccc ttt gag atg gac tat tat gta gta cgg cag ett tcc tgg gcg cat 628
 Pro Phe Glu Met Asp Tyr Tyr Val Val Arg Gln Leu Ser Trp Ala His
 105 110 115 120
 ggt cac gtg ctt tgt gcc tcc gtc aac tac ctt cgt acg gtc tcc ctg 676
 Gly His Val Leu Cys Ala Ser Val Asn Tyr Leu Arg Thr Val Ser Leu
 125 130 135

tac gtc tcc acc aac gct ctg ctg gcc atc gct att gac aga tac ctc Tyr Val Ser Thr Asn Ala Leu Leu Ala Ile Ala Ile Asp Arg Tyr Leu 140 145 150	724
gct att gtc cac cct ttg aaa cca cgg atg aat tat cag acc gct tcc Ala Ile Val His Pro Leu Lys Pro Arg Met Asn Tyr Gln Thr Ala Ser 155 160 165	772
ttc ctg atc gct ttg gtc tgg atg gtc tcc atc ctc atc gct gtc cca Phe Leu Ile Ala Leu Val Trp Met Val Ser Ile Leu Ile Ala Val Pro 170 175 180	820
tct gcc tac ttc acc aca gaa acc atc ctc gtt atc gtc aag aat caa Ser Ala Tyr Phe Thr Thr Glu Thr Ile Leu Val Ile Val Lys Asn Gln 185 190 195 200	868
gaa aaa atc ttc tgt ggt cag atc tgg tgg gtc gac cag cag ctc tac Glu Lys Ile Phe Cys Gly Gln Ile Trp Ser Val Asp Gln Val Leu Tyr 205 210 215	916
tac aaa tcc tac ttc ctc ttc gtc ttc ggg ctt gag ttc gtg ggt ccc Tyr Lys Ser Tyr Phe Leu Phe Val Phe Gly Leu Glu Phe Val Gly Pro 220 225 230	964
gtg gtc act atg acc ctg tgc tat gcc agg atc tcc caa gag ctc tgg Val Val Thr Met Thr Leu Cys Tyr Ala Arg Ile Ser Gln Glu Leu Trp 235 240 245	1012
ttc aag gct gta cct ggc ttc cag acg gag caa atc cgc aag cgg ctg Phe Lys Ala Val Pro Gly Phe Gln Thr Glu Gln Ile Arg Lys Arg Leu 250 255 260	1060
cgt tgc cgc cgc aag aca gtg cta ctg ctc atg ggc atc ctc aca gcc Arg Cys Arg Arg Lys Thr Val Leu Leu Leu Met Gly Ile Leu Thr Ala 265 270 275 280	1108
tac gtg ctg tgc tgg gcg ccg ttc tat ggc ttt acc ata gtg cga gac Tyr Val Leu Cys Trp Ala Pro Phe Tyr Gly Phe Thr Ile Val Arg Asp 285 290 295	1156
ttc ttc ccc acg gta gtt gtg aag gag aag cac tac ctc acc gcc ttc Phe Phe Pro Thr Val Val Val Lys Glu Lys His Tyr Leu Thr Ala Phe 300 305 310	1204
tac gtc gtg gag tgc att gcc atg agc aac agc atg atc aat act ata Tyr Val Val Glu Cys Ile Ala Met Ser Asn Ser Met Ile Asn Thr Ile 315 320 325	1252
tgc ttc gtg acg gtc aag aac aac acc atg aaa tac ttc aag aag atg Cys Phe Val Thr Val Lys Asn Asn Thr Met Lys Tyr Phe Lys Lys Met 330 335 340	1300
ctg cgg ctc cac tgg cgg ccc tct cac tac ggg agt aag tcc agc gct Leu Arg Leu His Trp Arg Pro Ser His Tyr Gly Ser Lys Ser Ser Ala 345 350 355 360	1348

gac ctc gac ctc aaa acc agc ggg gtg cct gcc act gaa gag gtg gat 1396
 Asp Leu Asp Leu Lys Thr Ser Gly Val Pro Ala Thr Glu Glu Val Asp 375
 365 370 375

tgt atc aga cta aag tagccttcag gtgtgcccc agggaaaaatt taacattcgg 1451
 Cys Ile Arg Leu Lys 380

tactcagtaa atcacacacc atcaaccact cacaagctac atggaaaagat acggctgtat 1511
 tcacgtctct ctgctctaat gtatcaggac gctctctatgt aataacatac agcacaactg 1571
 atgtctgtcat aacatcttag aagcagagaca caaataagtaa caagtgtatgt ggcactgaatg 1631
 ctctctgtcat caaacacacac caacaaatta ttcaaggaca agagctgaca tgtgagaatt 1691
 acctgctatg tgcaaaaaca agtlaccccc ccaaaaaatg atagaagcta tttggagtta 1751
 ttacagctcta tctatctatc tatctatcca tccatccatc catccatcca ggtcactaga 1811
 aagaagtccac aaatgactag ccagagtcatt gctacatatt ctttcattct gtatcttttc 1871
 tgacacagaac tgtcaaaggc aatagaataa agcacctaga catactagaa atgtaaggat 1931
 aactccatca ataggggagac caagggcctca taggaagagg gtccatatag tatactgact 1991
 ttccccatca cacaccagtt atctccttag atattctgta cttatctgca atgtgttaatt 2051
 ttcaaatgag gaaaaataag gggacaggct ttaccacaga tgtatcaaat cctcatcaagc 2111
 ccatagggaca aagatggggc gctcctgaca caagaaatgt atccagttct ggataaacttt 2171
 aatgccaaagc atttcagggc tctgggtgtc tggaggagaag ggacacagaa agagccagg 2231
 ttccagtggt caatgagttat aatctgtcca tttgctatga tttggacaat tttctagaac 2291
 atactccgag ttacaaaagg aactctactt gagatccaaa gatccgggta aaagtccctaa 2351
 cccccaggact catctctgtg tgtctccact gtaatgaaat ggaataaatg aaaacggatc 2411
 attaggaaca tcagcccgcc gaagtcattg tgtggatgtg atttccact cttcctttgt 2471
 gaagaatgag gtctgtgaaa gctcattaga gggagtttgg aatggagaaa cagctccaca 2531
 cttttcatcc ctctcttttg aatcggagac cactaaacgc atctttgaag tagcgtattct 2591
 atagtggaggc atcaaggctct cccgtccaca gagtgcacatc aagaaaaatc agtcaatgcc 2651
 cataccctca gcactccctgt ggtcttagac agtcttccca acaagcact ggtggaccgcc 2711
 aggactgaat tcaacttgtat tatatgtca tctactgaa atcagggttg atcaagttgg 2771
 ctagataggt attttcttcc tcttccaca ccccatatgt atccctccct taaatccagt 2831
 tactaaggaa gacctcttta aacacaggag aaccattatt cgttccagga cacaaataac 2891
 ctctccagta gacactgtac ccttcacatg tcaacagaat ttgctccctt cttgtattta 2951
 aacatcatcat cctaatttca tttagattta accagaaacc attcctgtaa atttcaatgt 3011
 gtttgtgata ccgactgtga aaaagcgtat gctgttatca tatggaataa ttaacataca 3071
 gaattgtaat cgtagtctcc aaaagggttcc ctactcctgt tgtattctat gtttatatgt 3131
 ttgatgtaaa tggagctgtg tagctgtcta agcagctcaa gctgaaatg agggaaatgtc 3191
 caatggtgtt cttagagcag ggccatctca ggctagcagc tggcctcagt ctgtgctctc 3251
 tcgggagtt gttcttaaat atgaattagc agcaaacat taaaaaaaaa aaaaaagggc 3311
 ggccgc 3317

<210> 4
 <211> 381
 <212> PRT
 <213> Mus musculus

<400> 4
 Met Gly Pro Gln Asn Arg Asn Thr Ser Phe Ala Pro Asp Leu Asn Pro
 1 5 10 15
 Pro Gln Asp His Val Ser Leu Asn Tyr Ser Tyr Gly Asp Leu
 20 25 30
 Pro Leu Gly Glu Asp Glu Asp Val Thr Lys Thr Gln Thr Phe Phe Ala
 35 40 45
 Ala Lys Ile Val Ile Gly Val Ala Leu Ala Gly Ile Met Leu Val Cys
 50 55 60
 Gly Ile Gly Asn Phe Val Phe Ile Ala Ala Leu Ala Arg Tyr Lys Lys
 65 70 75 80

Leu Arg Asn Leu Thr Asn Leu Leu Ile Ala Asn Leu Ala Ile Ser Asp
 85 90 95
 Phe Leu Val Ala Ile Val Cys Cys Pro Phe Glu Met Asp Tyr Tyr Val
 100 105 110
 Val Arg Gln Leu Ser Trp Ala His Gly His Val Leu Cys Ala Ser Val
 115 120 125
 Asn Tyr Leu Arg Thr Val Ser Leu Tyr Val Ser Thr Asn Ala Leu Leu
 130 135 140
 Ala Ile Ala Ile Asp Arg Tyr Leu Ala Ile Val His Pro Leu Lys Pro
 145 150 155 160
 Arg Met Asn Tyr Gln Thr Ala Ser Phe Leu Ile Ala Leu Val Trp Met
 165 170 175
 Val Ser Ile Leu Ile Ala Val Pro Ser Ala Tyr Phe Thr Thr Glu Thr
 180 185 190
 Ile Leu Val Ile Val Lys Asn Gln Glu Lys Ile Phe Cys Gly Gln Ile
 195 200 205
 Trp Ser Val Asp Gln Gln Leu Tyr Tyr Lys Ser Tyr Phe Leu Phe Val
 210 215 220
 Phe Gly Leu Glu Phe Val Gly Pro Val Val Thr Met Thr Leu Cys Tyr
 225 230 235 240
 Ala Arg Ile Ser Gln Glu Leu Trp Phe Lys Ala Val Pro Gly Phe Gln
 245 250 255
 Thr Glu Gln Ile Arg Lys Arg Leu Arg Cys Arg Arg Lys Thr Val Leu
 260 265 270
 Leu Leu Met Gly Ile Leu Thr Ala Tyr Val Leu Cys Trp Ala Pro Phe
 275 280 285
 Tyr Gly Phe Thr Ile Val Arg Asp Phe Phe Pro Thr Val Val Val Lys
 290 295 300
 Glu Lys His Tyr Leu Thr Ala Phe Tyr Val Val Glu Cys Ile Ala Met
 305 310 315 320
 Ser Asn Ser Met Ile Asn Trp Ile Cys Phe Val Thr Val Lys Asn Asn
 325 330 335
 Thr Met Lys Tyr Phe Lys Lys Met Leu Arg Leu His Trp Arg Pro Ser
 340 345 350
 His Tyr Gly Ser Lys Ser Ser Ala Asp Leu Asp Leu Lys Thr Ser Gly
 355 360 365
 Val Pro Ala Thr Glu Glu Val Asp Cys Ile Arg Leu Lys
 370 375 380

<210> 5
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer

<400> 5
 tgctgcttaa acctgggtcg g

21

<210> 6
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer

<400> 6
gggtgtgat ttactgagta ccg

23

<210> 7
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 7
cgccaccagg aagtcagaga tg

22


<210> 8
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer

<400> 8
gggaccaga acagaaacac ta

22

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



SCANNED, # 1A

Application deficiencies were found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

There are 100 sheets of specification.

☐ Scanned copy is best available.